

THE BAILEY LEMMA AND KOSTKA POLYNOMIALS

S. OLE WARNAAR

ABSTRACT. Using the theory of Kostka polynomials, we prove an A_{n-1} version of Bailey's lemma at integral level. Exploiting a new, conjectural expansion for Kostka numbers, this is then generalized to fractional levels, leading to a new expression for admissible characters of $A_{n-1}^{(1)}$ and to identities for A-type branching functions.

1. INTRODUCTION

Let $\alpha = (\alpha_L)_{L \geq 0}$ and $\beta = (\beta_L)_{L \geq 0}$ be sequences and $(a; q)_r = (a)_r = \prod_{k=0}^{r-1} (1 - aq^k)$ a q -shifted factorial. Then (α, β) is called a Bailey pair relative to a if

$$(1.1) \quad \beta_L = \sum_{r=0}^L \frac{\alpha_r}{(q)_{L-r} (aq)_{L+r}}.$$

Similarly, a pair of sequences (γ, δ) is called a conjugate Bailey pair relative to a if

$$(1.2) \quad \gamma_L = \sum_{r=L}^{\infty} \frac{\delta_r}{(q)_{r-L} (aq)_{r+L}}.$$

By a simple interchange of sums it follows that if (α, β) and (γ, δ) are a Bailey pair and conjugate Bailey pair relative to a , then

$$(1.3) \quad \sum_{L=0}^{\infty} \alpha_L \gamma_L = \sum_{L=0}^{\infty} \beta_L \delta_L,$$

provided of course that all sums converge.

Bailey used (1.3) for proving identities of the Rogers–Ramanujan type [7]. First he showed that a limiting case of the q -Gauss sum [13, Eq. (II.8)] provides the following conjugate Bailey pair relative to a :

$$(1.4) \quad \gamma_L = \frac{a^L q^{L^2}}{(aq)_{\infty}} \quad \text{and} \quad \delta_L = a^L q^{L^2} \quad (|q| < 1).$$

Substituted into (1.3) this gives

$$(1.5) \quad \frac{1}{(aq)_{\infty}} \sum_{L=0}^{\infty} a^L q^{L^2} \alpha_L = \sum_{L=0}^{\infty} a^L q^{L^2} \beta_L.$$

To obtain the Rogers–Ramanujan identities, Bailey now required [6]

$$(1.6) \quad \alpha_L = \frac{1 - aq^{2L}}{1 - a} \frac{(-a)^L q^{L(3L-1)/2} (a)_L}{(q)_L} \quad \text{and} \quad \beta_L = \frac{1}{(q)_L},$$

which follows from a limit of Rogers' q -analogue of Dougall's ${}_5F_4$ sum [13, Eq. (II.21)]. Inserting (1.6) into (1.5) yields the Rogers–Selberg identity. Taking $a = 1$ and $a = q$ and performing the

2000 *Mathematics Subject Classification*. Primary 05A19, 05E05, 11B65; Secondary 17B65, 33D15.
Work supported by the Australian Research Council.

sum on the left using Jacobi's triple product identity [13, Eq. (II.28)] results in the Rogers–Ramanujan identities [39]

$$(1.7) \quad \sum_{n=0}^{\infty} \frac{q^{n^2}}{(q)_n} = \frac{1}{(q, q^4; q^5)_{\infty}} \quad \text{and} \quad \sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q)_n} = \frac{1}{(q^2, q^3; q^5)_{\infty}},$$

where $(a_1, \dots, a_k; q)_r = (a_1; q)_r \cdots (a_k; q)_r$ and $|q| < 1$.

About a decade ago Milne and Lilly generalized the notion of a Bailey pair to higher dimensions [36]. (For a recent review see [35].) They defined, for example, an A_{n-1} Bailey pair relative to a by

$$\beta_L = \sum_{\substack{r_i=0 \\ 1 \leq i \leq n-1}}^{L_i} M_{L,r} \alpha_r,$$

where $L, r \in \mathbb{Z}_+^{n-1}$ and

$$M_{L,r} = \prod_{j=1}^{n-1} \left[(aqx_j/x_{n-1})_{L_j+|r|} \prod_{k=1}^{n-1} (q^{r_j-r_k+1}x_j/x_k)_{L_j-r_k} \right]^{-1},$$

with x_1, \dots, x_{n-1} indeterminates and $|r| = r_1 + \cdots + r_{n-1}$. From the point of view of A_{n-1} basic hypergeometric series the Milne–Lilly definition of an A_{n-1} Bailey pair is not only natural but, as the papers [34, 36, 37] attest, has also been very fruitful. However, from the point of view of symmetric functions a rather different definition of an A_{n-1} Bailey pair (and conjugate Bailey pair) seems natural. To motivate this, let us assume that $a = q^\ell$ in (1.1) with ℓ a nonnegative integer. Then

$$(1.8) \quad \beta_L = \sum_{r=0}^L \frac{\alpha_r}{(aq)_{2L}} \begin{bmatrix} 2L + \ell \\ L - r \end{bmatrix}_q,$$

where $\begin{bmatrix} n \\ m \end{bmatrix}_q$ is a q -binomial coefficient, defined as

$$(1.9) \quad \begin{bmatrix} n \\ m \end{bmatrix}_q = \begin{cases} \frac{(q)_n}{(q)_m (q)_{n-m}} & \text{for } 0 \leq m \leq n \\ 0 & \text{otherwise.} \end{cases}$$

For λ, μ partitions and η a composition let $K_{\lambda\eta}$ and $K_{\lambda\mu}(q)$ be a Kostka number and Kostka polynomial, respectively. (For their definition and for notation concerning partitions and compositions, see Section 2.1.) From [32, Ch. I, §6, Ex. 2] it follows that for $\lambda = (\lambda_1, \lambda_2)$ and $\eta = (\eta_1, \eta_2)$ there holds

$$(1.10) \quad K_{\lambda\eta} = \chi(\lambda_2 \leq \eta_1 \leq \lambda_1) \chi(|\lambda| = |\eta|),$$

where $\chi(\text{true}) = 1$ and $\chi(\text{false}) = 0$. Also, from [32, Ch. III, §6, Ex. 2]

$$K_{(1^i-2^j 2^j), (1^i)}(q) = \begin{bmatrix} i \\ j \end{bmatrix}_q - \begin{bmatrix} i \\ j-1 \end{bmatrix}_q.$$

Combining these formulae leads to the expansion

$$\begin{aligned}
(1.11) \quad \begin{bmatrix} L \\ r \end{bmatrix}_q &= \sum_{k=0}^{\min\{r, L-r\}} \left\{ \begin{bmatrix} L \\ k \end{bmatrix}_q - \begin{bmatrix} L \\ k-1 \end{bmatrix}_q \right\} \\
&= \sum_{k=0}^{\lfloor L/2 \rfloor} K_{(L-k, k), (r, L-r)} K_{(1^{L-2k} 2^k), (1^L)}(q) \\
&= \sum_{\mu \vdash L} K_{\mu, (r, L-r)} K_{\mu', (1^L)}(q),
\end{aligned}$$

where in the last step we have used that $K_{\mu, (r, L-r)} = 0$ if $l(\mu) > 2$. Substituted into (1.8) this results in

$$\begin{aligned}
(1.12) \quad \beta_L &= \sum_{r=0}^L \frac{\alpha_r}{(aq)_{2L}} \sum_{\mu \vdash 2L+\ell} K_{\mu, (L-r, L+r+\ell)} K_{\mu', (1^{2L+\ell})}(q) \\
&= (q)_\ell \sum_{r=0}^{(|\eta|-\ell)/2} \frac{\alpha_r}{(q)_{m(\eta)}} \sum_{\mu \vdash |\eta|} K_{\mu, \frac{|\eta|}{2}(1^2) - k - \ell \bar{\Lambda}_1} K_{\mu', \eta}(q)
\end{aligned}$$

where $\eta = (1^{2L+\ell})$, $(q)_{m(\eta)} = (q)_{2L+\ell}$, $\bar{\Lambda}_1 = (1/2, -1/2)$ and $k = (r, -r)$. This suggests that a natural multivariable or A_{n-1} generalization of a Bailey pair is obtained by letting η be a partition of at most $n-1$ different parts and by replacing the composition $|\eta|(1^2)/2 - k + \ell \bar{\Lambda}_1$ by an appropriate composition of n parts. Such a generalization was introduced in [44], and various conjectures were made concerning A_{n-1} Bailey pairs and conjugate Bailey pairs. Further progress was made in [5], where it was shown that the A_2 generalization of (1.12) leads to A_2 versions of the Rogers–Ramanujan identities (1.7) such as

$$\sum_{n_1, n_2=0}^{\infty} \frac{q^{n_1^2 - n_1 n_2 + n_2^2}}{(q)_{n_1}} \begin{bmatrix} 2n_1 \\ n_2 \end{bmatrix}_q = \frac{1}{(q, q, q^3, q^4, q^6, q^6; q^7)_\infty}.$$

In this paper we will study the more general A_{n-1} generalization of (1.12) and prove the conjectured conjugate Bailey pairs of [44]. Unlike in [5, 42, 45, 46, 52], where conjugate Bailey pairs are proven using summation and transformation formulas for basic hypergeometric series, our approach here will be to employ the theory of Kostka polynomials. Even in the classical or A_1 case this is new. Having proven the conjectures of [44], we will utilize a conjectural expansion for Kostka numbers to derive much more general conjugate Bailey pairs in terms of fractional-level string functions and configuration sums of $A_{n-1}^{(1)}$. Not only will this give rise to new expressions for admissible characters of $A_{n-1}^{(1)}$, but also to q -series identities for unitary as well as nonunitary A-type branching functions.

The outline of this paper is as follows. The next section contains an introduction to symmetric functions, Kostka polynomials and supernomial coefficients, and concludes with a conjectured expansion of the Kostka numbers in terms of antisymmetric supernomials. In Section 3 we define an A_{n-1} analogue of Bailey's lemma and prove a generalization of the conjugate Bailey pair (1.4), settling a conjecture of [44]. Section 4 provides a bosonic reformulation of the results of Section 3, which is exploited in the subsequent section to yield very general conjugate Bailey pairs in terms of $A_{n-1}^{(1)}$ string functions and configuration sums. Finally, in Section 6, we use these results to find many new A-type q -series identities.

2. SYMMETRIC FUNCTIONS, KOSTKA POLYNOMIALS AND SUPERNOMIAL COEFFICIENTS

2.1. Introduction. This section contains a brief introduction to symmetric functions and Kostka polynomials. For further details, see [10, 12, 32, 33].

A composition $\lambda = (\lambda_1, \lambda_2, \dots)$ is a sequence (finite or infinite) of nonnegative integers with finitely many $\lambda_i \geq 0$ unequal to zero. Two compositions that only differ in their tails of zeros are identified, and, for example, we will not distinguish between $(1, 0, 2, 3, 0, 0)$ and $(1, 0, 2, 3)$. Even when writing $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}_+^n$ for a composition λ we do give meaning to λ_{n+k} for $k \geq 1$ (namely the obvious $\lambda_{n+k} = 0$). The weight of a composition λ , denoted $|\lambda|$, is the sum of its (nonzero) components. The largest and smallest components of a composition λ will be denoted by $\max(\lambda)$ and $\min(\lambda)$. Also for noninteger sequences μ will we write $|\mu|$ for the sum of its components.

A partition λ is a composition whose components are weakly decreasing. The nonzero components of a partition λ are called its parts and the number of parts, denoted $l(\lambda)$, its length. We say that λ is a partition of k , denoted $\lambda \vdash k$ if $|\lambda| = k$.

The Ferrers graph of a partition λ is obtained by drawing $l(\lambda)$ left-aligned rows of dots, with the i th row containing λ_i dots. A partition is called rectangular if all its parts are equal, i.e., if its Ferrers graph has rectangular shape. The conjugate of a partition λ , denoted λ' , is obtained by transposing the Ferrers graph of λ . For λ a partition,

$$n(\lambda) = \sum_{i=1}^{l(\lambda)} (i-1)\lambda_i = \sum_{i=1}^{\lambda_1} \binom{\lambda'_i}{2}.$$

The unique partition $(0, 0, \dots)$ of 0 is denoted by \emptyset , the set of all partitions of k by \mathcal{P}_k and the set of all partitions by \mathcal{P} . We define the usual dominance (partial) order on \mathcal{P}_k by writing $\lambda \geq \mu$ for $\lambda, \mu \in \mathcal{P}_k$ iff $\lambda_1 + \dots + \lambda_i \geq \mu_1 + \dots + \mu_i$ for all $i \geq 1$. Note that $\lambda \geq \mu \Leftrightarrow \mu' \geq \lambda'$ and that $\lambda \geq \mu$ implies $\lambda_1 \geq \mu_1$ as well as $l(\lambda) \leq l(\mu)$.

Frequently $\lambda \in \mathcal{P}$ will be written as $\lambda = (1^{m_1} 2^{m_2} \dots)$ where $m_j = \lambda'_j - \lambda'_{j+1}$ is the multiplicity of the part j . The associated composition $(m_1, m_2, \dots, m_{\lambda_1})$ is denoted by $m(\lambda)$. Using the multiplicities, the union of the partitions $\lambda = (1^{m_1} 2^{m_2} \dots)$ and $\mu = (1^{n_1} 2^{n_2} \dots)$ is given by $\lambda \cup \mu = (1^{m_1+n_1} 2^{m_2+n_2} \dots)$. For λ a partition such that $\lambda_1 \leq a$

$$(2.1) \quad n((a^b) \cup \lambda) = a \binom{b}{2} + b|\lambda| + n(\lambda).$$

One can also define the sum of partitions but, more generally, we will simply assume the standard addition and subtraction for sequences in \mathbb{R}^n . For two partitions λ and μ we write $\mu \subseteq \lambda$ if $\mu_i \leq \lambda_i$ for all i , and $\nu = \lambda \cap \mu$ if $\nu_i = \min\{\lambda_i, \mu_i\}$ for all i .

Let S_n be the symmetric group, i.e., the group with elements given by the permutations of $(1, \dots, n)$, and with multiplication \circ given by the usual composition of permutations. Given a permutation $\sigma \in S_n$ let $\ell(\sigma)$ be its length (= the minimal number of ‘‘adjacent transpositions’’ required to obtain σ from $(1, 2, \dots, n)$) and $\epsilon(\sigma) = (-1)^{\ell(\sigma)}$ its signature. Clearly $\epsilon(\sigma \circ \tau) = \epsilon(\sigma)\epsilon(\tau)$ and (thus) $\epsilon(\sigma) = \epsilon(\sigma^{-1})$. The unique permutation of maximal length (= $n(n-1)/2$) will be denoted by π . (Incidentally, π is the partition $(n, \dots, 2, 1)$ which also has length $l(\pi) = n$ in the sense of partitions.) Let $k = (k_1, \dots, k_n)$. Then $\sigma \in S_n$ acts on k by permuting its components; $\sigma(k) = (k_{\sigma_1}, \dots, k_{\sigma_n})$. For λ a partition such that $l(\lambda) \leq n$, S_n^λ denotes the subgroup of S_n consisting of permutations that leave λ invariant, i.e., $\sigma(\lambda) = \lambda$ for $\sigma \in S_n^\lambda$.

Let $x^\lambda = x_1^{\lambda_1} \dots x_n^{\lambda_n}$. For λ a partition such that $l(\lambda) \leq n$ the monomial symmetric function in n variables is defined as

$$m_\lambda(x) = \sum_{\sigma \in S_n / S_n^\lambda} x^{\sigma(\lambda)}.$$

If $l(\lambda) > n$, $m_\lambda(x) = 0$. To define the complete symmetric function h_λ , set $h_r(x) = 0$ for $r < 0$,

$$h_r(x) = \sum_{\mu \vdash r} m_\mu(x) = \sum_{1 \leq i_1 \leq \dots \leq i_r \leq n} x_{i_1} \cdots x_{i_r}.$$

for $r \geq 0$, and $h_\lambda(x) = h_{\lambda_1}(x)h_{\lambda_2}(x)\cdots$ for λ a composition. In much the same way the elementary symmetric function e_λ is defined by $e_0(x) = 1$,

$$e_r(x) = m_{(1^r)}(x) = \sum_{1 \leq i_1 < \cdots < i_r \leq n} x_{i_1} \cdots x_{i_r}$$

for $r \geq 1$, and $e_\lambda(x) = e_{\lambda_1}(x)e_{\lambda_2}(x)\cdots$ for λ a composition. Since $e_r(x) = 0$ for $r > n$ we have $e_\lambda(x) = 0$ if $\max(\lambda) > n$ ($\lambda_1 > n$ for $\lambda \in \mathcal{P}$). Clearly, $e_\lambda(x) = e_{\sigma(\lambda)}(x)$ and $h_\lambda(x) = h_{\sigma(\lambda)}(x)$.

For $\lambda \in \mathcal{P}$ such that $l(\lambda) \leq n$ let

$$(2.2) \quad a_\lambda(x) = \sum_{\sigma \in S_n} \epsilon(\sigma) x^{\sigma(\lambda)}.$$

Then the Schur function $s_\lambda(x)$ is defined as

$$s_\lambda(x) = \frac{a_{\lambda+\delta}(x)}{a_\delta(x)},$$

where $\delta = (n-1, n-2, \dots, 1, 0)$. The denominator on the right is known as the Vandermonde determinant;

$$(2.3) \quad a_\delta(x) = \prod_{1 \leq i < j \leq n} (x_i - x_j).$$

Finally we introduce the Hall-Littlewood symmetric function as the following q -analogue of the monomial symmetric function:

$$P_\lambda(x; q) = \sum_{\sigma \in S_n / S_n^\lambda} x^{\sigma(\lambda)} \prod_{\lambda_i > \lambda_j} \frac{x_{\sigma_i} - qx_{\sigma_j}}{x_{\sigma_i} - x_{\sigma_j}}$$

for $\lambda \in \mathcal{P}$ such that $l(\lambda) \leq n$, and $P_\lambda(x; q) = 0$ for $\lambda \in \mathcal{P}$ such that $l(\lambda) > n$. Besides the obvious $P_\lambda(x; 1) = m_\lambda(x)$ one also has $P_\lambda(x; 0) = s_\lambda(x)$.

For μ a composition and $\lambda \vdash |\mu|$, the Kostka number $K_{\lambda\mu}$ is defined by

$$(2.4) \quad h_\mu(x) = \sum_{\lambda \vdash |\mu|} K_{\lambda\mu} s_\lambda(x).$$

More generally, $K_{\lambda\mu} = 0$ if λ is not a partition, or μ not a composition, or $|\mu| \neq |\lambda|$. From $h_\mu(x) = h_{\sigma(\mu)}(x)$ it follows that

$$(2.5) \quad K_{\lambda\mu} = K_{\lambda, \sigma(\mu)}.$$

Further simple properties of the Kostka numbers are

$$(2.6) \quad K_{\lambda\mu} \neq 0 \text{ iff } \lambda \geq \mu$$

for $\lambda \in \mathcal{P}$ and $\mu \vdash |\lambda|$, and

$$(2.7) \quad K_{\lambda\mu} = K_{(a^n)+\lambda, (a^n)+\mu}$$

for $l(\lambda) \leq n$, $\mu \in \mathbb{Z}_+^n$ and $a \geq -\lambda_n$.

Occurrences of the Kostka numbers similar to (2.4) are given by

$$(2.8) \quad e_\mu(x) = \sum_{\lambda \vdash |\mu|} K_{\lambda'\mu} s_\lambda(x)$$

for μ a composition, and

$$(2.9) \quad s_\lambda(x) = \sum_{\mu \vdash |\lambda|} K_{\lambda\mu} m_\mu(x)$$

for $\lambda \in \mathcal{P}$. Note that both sides of (2.8) vanish if $\max(\mu) > n$, and both sides of (2.9) vanish if $l(\lambda) > n$.

A useful formula involving the Kostka numbers follows from the Jacobi-Trudi identity

$$s_\mu(x) = \det(h_{\mu_i+j-i}(x))_{1 \leq i, j \leq n} = \sum_{\sigma \in S_n} \epsilon(\sigma) h_{\sigma(\mu+\delta)-\delta}.$$

Substituting (2.4) and interchanging sums gives

$$s_\mu(x) = \sum_{\lambda \vdash |\mu|} s_\lambda(x) \sum_{\sigma \in S_n} \epsilon(\sigma) K_{\lambda, \sigma(\mu+\delta)-\delta},$$

which implies that

$$(2.10) \quad \sum_{\sigma \in S_n} \epsilon(\sigma) K_{\lambda, \sigma(\mu+\delta)-\delta} = \delta_{\lambda, \mu}$$

for $\lambda, \mu \in \mathcal{P}$ such that $l(\mu) \leq n$ and $\delta_{a,b} = \chi(a = b)$. Note that (2.10) also holds for $\mu = (\mu_1, \mu_2, \dots, \mu_n)$ such that $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n$ and $\mu_n < 0$, since $\sigma(\mu + \delta) - \delta$ is not a composition for such a μ .

For $\lambda, \mu \in \mathcal{P}$ the Kostka polynomial (or Kostka–Foulkes polynomial) $K_{\lambda\mu}(q)$ is defined by

$$(2.11) \quad s_\lambda(x) = \sum_{\mu \vdash |\lambda|} K_{\lambda\mu}(q) P_\mu(x; q)$$

and $K_{\lambda\mu}(q) = 0$ if $|\lambda| \neq |\mu|$. We extend this to compositions μ through $K_{\lambda\mu}(q) = K_{\lambda, \sigma(\mu)}(q)$ and to more general sequences λ and μ by $K_{\lambda\mu} = 0$ if λ is not a partition, or μ is not a composition. Since $P_\lambda(x; 1) = m_\lambda(x)$ it follows that $K_{\lambda\mu}(1) = K_{\lambda\mu}$.

Let $\lambda, \mu \in \mathcal{P}$ such that r is rectangular and $\lambda, \mu \subseteq r$, and denote by $\tilde{\lambda}_r$ ($\tilde{\mu}_r$) the “complement of λ (μ) with respect to r ”. That is, for $r = (a^b)$,

$$\tilde{\lambda}_r = (\underbrace{a, \dots, a}_{b-l(\lambda)}, a - \lambda_{l(\lambda)}, \dots, a - \lambda_1).$$

Then the following duality holds [28]:

$$(2.12) \quad K_{\lambda\mu}(q) = K_{\tilde{\lambda}_r, \tilde{\mu}_r}(q).$$

For example, by taking $r = (4^3)$,

$$K_{(3,1),(2,1,1)}(q) = K_{(4,3,1),(3,3,2)}(q) = q + q^2.$$

Note that (2.12) implies a q -analogue of (2.7) as well as

$$(2.13) \quad K_{\lambda\mu}(q) = K_{(n^a) \cup \lambda, (n^a) \cup \mu}(q)$$

for $\lambda, \mu \in \mathcal{P}$ such that $\lambda_1, \mu_1 \leq n$ and $a \geq 0$.

2.2. \mathbf{A}_{n-1} supernomial coefficients. We now come to perhaps the two most important definitions, that of the completely symmetric and completely antisymmetric \mathbf{A}_{n-1} supernomials [15, 43, 44]. Although all that is needed for their definition is at our disposal, we shall somewhat digress and motivate the supernomials from the point of view of generalized Kostka polynomials. In [26, 44, 47, 48, 49, 50] generalizations of the Kostka polynomials $K_{\lambda\mu}(q)$ were introduced wherein the composition μ in the ordinary Kostka polynomials is replaced by a (finite) sequence $R = (R_1, R_2, \dots)$ of rectangular partitions R_i , such that

$$(2.14) \quad K_{\lambda R}(q) = K_{\lambda\mu}(q) \quad \text{if } R_i = (\mu_i) \text{ for all } i.$$

(For $R = (R_1)$, $K_{\lambda R}(q) = \delta_{\lambda, R_1}$ and *not* $K_{\lambda R}(q) = K_{\lambda R_1}(q)$.) Whereas $K_{\lambda\mu}$ is the cardinality of the set of (semi-standard) Young tableaux of shape λ and content (or weight) μ [32], $K_{\lambda R}(1)$ is the cardinality of the set of Littlewood–Richardson tableaux of shape λ and content R [30]. This implies that $K_{\lambda R}(q) = 0$ if $|\lambda| \neq \sum_i |R_i| = |R|$. Equivalently, the generalized Kostka

polynomials may be viewed as q -analogues of generalized Littlewood–Richardson coefficients. Recalling the definition of the Littlewood–Richardson coefficients $c_{\lambda\mu}^\nu$ as

$$(2.15) \quad s_\lambda(x)s_\mu(x) = \sum_{\nu \in \mathcal{P}} c_{\lambda\mu}^\nu s_\nu(x)$$

for $\lambda, \mu \in \mathcal{P}$ (hence $c_{\lambda\mu} = 0$ if $|\nu| \neq |\lambda| + |\mu|$), there holds

$$(2.16) \quad K_{\lambda R}(1) = \sum_{\nu^{(1)}, \dots, \nu^{(k-2)} \in \mathcal{P}} c_{\nu^{(0)}R_2}^{\nu^{(1)}} c_{\nu^{(1)}R_3}^{\nu^{(2)}} \cdots c_{\nu^{(k-2)}R_k}^{\nu^{(k-1)}},$$

with $R = (R_1, \dots, R_k)$, $\nu^{(0)} = R_1$ and $\nu^{(k-1)} = \lambda$. For $k = 2$ this yields $K_{\lambda, (R_1, R_2)}(1) = c_{R_1 R_2}^\lambda$.

Now let $\lambda \in \mathbb{Z}_+^n$. Then the A_{n-1} supernomials are defined in terms of the generalized Kostka polynomials as

$$(2.17) \quad S_{\lambda R}(q) = \sum_{\nu \vdash |\lambda|} K_{\nu\lambda} K_{\nu'R^*}(q),$$

where $R^* = (R'_1, R'_2, \dots)$. From the duality [26, Prop. 12], [44, Thm. 7.1]

$$(2.18) \quad K_{\lambda R}(q) = q^{\sum_{i < j} |R_i \cap R_j|} K_{\lambda'R^*}(1/q)$$

it follows that $S_{\lambda R}(1) = \sum_{\nu \vdash |\lambda|} K_{\nu\lambda} K_{\nu R}(1)$. Multiplying this by $m_\lambda(x)$, summing over λ and using (2.9), (2.15) and (2.16), one finds

$$(2.19) \quad s_{R_1}(x)s_{R_2}(x) \cdots = \sum_{\lambda \vdash |R|} S_{\lambda R}(1)m_\lambda(x) = \sum_{\substack{\lambda \in \mathbb{Z}_+^n \\ |\lambda| = |R|}} S_{\lambda R}(1)x^\lambda.$$

The $S_{\lambda R}(1)$ may thus be viewed as a generalized multinomial coefficient.

In the following we need two special cases of the A_{n-1} supernomials. One, the completely antisymmetric A_{n-1} supernomial (antisymmetric supernomial for short) $\mathcal{A}_{\lambda\mu}(q)$, arises when $\lambda \in \mathbb{Z}_+^n$ and $R_i = (1^{\mu_i})$ for all i . Since $R_i^* = R'_i = (\mu_i)$, (2.14) and (2.17) imply

$$(2.20) \quad \mathcal{A}_{\lambda\mu}(q) := S_{\lambda((1^{\mu_1}), (1^{\mu_2}), \dots)}(q) = \sum_{\nu \vdash |\lambda|} K_{\nu\lambda} K_{\nu'\mu}(q).$$

By (2.8) and (2.11) it thus follows that $e_\mu(x) = \sum_{\lambda \vdash |\mu|} \mathcal{A}_{\lambda\mu}(q) P_\lambda(x; q)$. Since $P_\lambda(x; 1) = m_\lambda(x)$ and $e_\mu(x) = s_{(1^{\mu_1})}(x)s_{(1^{\mu_2})}(x) \cdots$, this is in accordance with (2.19).

In similar fashion the completely symmetric A_{n-1} supernomial $\mathcal{S}_{\lambda\mu}(q)$ results when $\lambda \in \mathbb{Z}_+^n$, $R_i = (\mu_i)$ for all i , and $q \rightarrow 1/q$;

$$(2.21) \quad \mathcal{S}_{\lambda\mu}(q) := q^{\sum_{i < j} \min\{\mu_i, \mu_j\}} S_{\lambda((\mu_1), (\mu_2), \dots)}(1/q) = \sum_{\nu \vdash |\lambda|} K_{\nu\lambda} K_{\nu\mu}(q),$$

where the expression on the right follows from (2.14), (2.17) and (2.18). Observe that when μ is a partition $\sum_{i < j} \min\{\mu_i, \mu_j\} = n(\mu)$. By (2.4) and (2.11) it follows that $h_\mu(x) = \sum_{\lambda \vdash |\mu|} \mathcal{S}_{\lambda\mu}(q) P_\lambda(x; q)$. Since $P_\lambda(x; 1) = m_\lambda(x)$ and $h_\mu(x) = s_{(\mu_1)}(x)s_{(\mu_2)}(x) \cdots$, this is again in agreement with (2.19).

The antisymmetric supernomial of (2.20) will play a particularly important role throughout the paper and it will be convenient to introduce the further notation

$$(2.22) \quad \begin{bmatrix} L \\ \lambda \end{bmatrix} = \mathcal{A}_{\lambda\mu}(q) = \sum_{\nu \vdash |\lambda|} K_{\nu\lambda} K_{\nu'\mu}(q),$$

for $\lambda \in \mathbb{Z}_+^n$, $\mu \in \mathcal{P}_\lambda$ such that $\mu_1 \leq n-1$ and $L = m(\mu) \in \mathbb{Z}_+^{n-1}$. When $\lambda \in \mathbb{Z}^n$ with one or more components being negative we set $\begin{bmatrix} L \\ \lambda \end{bmatrix} = 0$. It is in fact not a restriction to assume that μ is a partition with largest part at most $n-1$. To see this, assume that $\lambda, \mu \in \mathcal{P}$ (with $l(\lambda) \leq n$), which is harmless since $\mathcal{A}_{\lambda\mu}(q) = \mathcal{A}_{\sigma(\lambda), \tau(\mu)}(q)$. Then the summand in

(2.20) is nonzero iff $\nu \geq \lambda$ and $\nu' \geq \mu$ implying that $\mathcal{A}_{\lambda\mu}(q)$ is nonzero iff $\mu' \geq \lambda$. One may therefore certainly assume that $\mu_1 \leq l(\lambda) \leq n$. But from (2.7) and (2.13) it is easily seen that $\mathcal{A}_{\lambda+(1^n), (n)\cup\mu}(q) = \mathcal{A}_{\lambda\mu}(q)$, so that we may actually assume $\mu_1 \leq n-1$ in $\mathcal{A}_{\lambda\mu}(q)$, leading naturally to (2.22).

In the case of A_1 , $\lambda = (\lambda_1, \lambda_2)$, $\mu = (1^{|\lambda|})$ and $L = m(\mu) = (|\lambda|)$, and from the extremes of (1.11) it follows that

$$(2.23) \quad \begin{bmatrix} (\lambda_1 + \lambda_2) \\ (\lambda_1, \lambda_2) \end{bmatrix} = \begin{bmatrix} \lambda_1 + \lambda_2 \\ \lambda_1 \end{bmatrix}_q,$$

with on the right the classical q -binomial coefficient (1.9). Two important symmetries of the antisymmetric supernomials needed subsequently are

$$(2.24) \quad \begin{bmatrix} L \\ \lambda \end{bmatrix} = \begin{bmatrix} L \\ \sigma(\lambda) \end{bmatrix}$$

and

$$(2.25) \quad \begin{bmatrix} (L_1, \dots, L_{n-1}) \\ \lambda \end{bmatrix} = \begin{bmatrix} (L_{n-1}, \dots, L_1) \\ (|L|^n) - \lambda \end{bmatrix}.$$

We will also use that the Kostka polynomials may be expressed in terms of antisymmetric supernomials as

$$(2.26) \quad K_{\lambda\mu}(q) = \sum_{\sigma \in S_n} \epsilon(\sigma) \begin{bmatrix} m(\mu) \\ \sigma(\lambda' + \delta) - \delta \end{bmatrix}.$$

Since (2.24) follows from (2.5) and (2.22), and (2.26) from (2.10) and (2.22), we only need to prove (2.25).

Proof of (2.25). By (2.12), $\tilde{\lambda}'_{r'} = (\tilde{\lambda}_r)'$ and $|\nu| + |\tilde{\nu}_r| = |r|$,

$$\begin{aligned} \mathcal{A}_{\lambda\mu'}(q) &= \sum_{\nu \vdash |\lambda|} K_{\nu\lambda} K_{\nu'\mu'}(q) = \sum_{\nu \vdash |\lambda|} K_{\tilde{\nu}_r, \tilde{\lambda}_r} K_{\tilde{\nu}'_{r'}, \tilde{\mu}'_{r'}}(q) \\ &= \sum_{\nu \vdash |\lambda|} K_{\tilde{\nu}_r, \tilde{\lambda}_r} K_{(\tilde{\nu}_r)', (\tilde{\mu}_r)'}(q) = \sum_{\tilde{\nu}_r \vdash |\tilde{\lambda}_r|} K_{\tilde{\nu}_r, \tilde{\lambda}_r} K_{(\tilde{\nu}_r)', (\tilde{\mu}_r)'}(q) = \mathcal{A}_{\tilde{\lambda}_r, (\tilde{\mu}_r)'}(q) \end{aligned}$$

for r a rectangular partition such that $\lambda, \mu \subseteq r$. Note that the summand in the second expression of the first line is nonzero for $\lambda \leq \nu \leq \mu$ only. This implies that if $\mu, \lambda \subseteq r$ then also $\nu \subseteq r$, and thus permits the use of (2.12) resulting in the third expression of the top-line.

Now specialize $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathcal{P}$, $\mu' = (1^{L_1}, \dots, (n-1)^{L_{n-1}})$ and $r = (\mu'_1) = (|L|^n)$, where $L = m(\mu')$. Then $\tilde{\lambda}_r = (|L|^n) - \pi(\lambda)$ and $(\tilde{\mu}_r)_i = L_1 + \dots + L_{n-i}$ so that $(\tilde{\mu}_r)' = (1^{L_{n-1}}, \dots, (n-1)^{L_1})$. Recalling (2.22), (2.25) now follows for $\lambda \in \mathcal{P}$ with the bottom-entry of the right side being substituted by $(|L|^n) - \pi(\lambda)$. Thanks to (2.24) this may be replaced by $(|L|^n) - \lambda$ with λ a composition. \square

2.3. A conjecture. To conclude this section we present a conjectured expansion of the Kostka numbers in terms of antisymmetric supernomials. Admitting this conjecture leads to an orthogonality relation between Kostka polynomials and antisymmetric supernomials (Corollary 2.2 below), which will be crucial in Section 5.

Before we can state the conjecture some more notation is needed. Denote the set $\{1, \dots, k\}$ by $[k]$. Let ε_i for $i \in [n]$ be the canonical basis vectors in \mathbb{R}^n with standard inner product $(\varepsilon_i | \varepsilon_j) = \delta_{i,j}$. Occasionally we will abbreviate $(v|v) = \|v\|^2$. Expressed in terms of the ε_i , the simple roots and fundamental weights α_i and $\bar{\Lambda}_i$ ($i \in [n-1]$) of A_{n-1} are given by $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$ and $\bar{\Lambda}_i = \varepsilon_1 + \dots + \varepsilon_i - \frac{i}{n}(1^n)$, respectively. (The notation Λ_i will be reserved for the fundamental weights of $A_{n-1}^{(1)}$.) The Weyl vector ρ is given by the sum of the fundamental weights; $\rho = \frac{1}{2}(n-1, n-3, \dots, 1-n) = \delta - (n-1)(1^n)/2$. The A_{n-1} weight and root lattices P

and $Q \subset P$ are the integral span of the fundamental weights and the integral span of the simple roots, respectively. $P_+ \subset P$ denotes the set of dominant weights, i.e., those weights $\lambda \in P$ for which $(\alpha_i|\lambda) \geq 0$ for all i . Specifically, for $\lambda \in P$ we have $\lambda \in \mathbb{Z}^n/n$, $|\lambda| = 0$ and $\lambda_i - \lambda_{i+1} \in \mathbb{Z}$. For $\lambda \in P_+$ the third condition needs to be sharpened to $\lambda_i - \lambda_{i+1} \in \mathbb{Z}_+$, and for $\lambda \in Q$ the first condition needs to be replaced by $\lambda \in \mathbb{Z}^n$. The weight and root lattices are invariant under the action of S_n , and for $\lambda \in P$ and $\sigma \in S_n$, $\sigma(\lambda) - \lambda \in Q$. This last fact easily follows from $\sigma^{(j)}(\bar{\Lambda}_i) = \bar{\Lambda}_i - \alpha_i \delta_{i,j}$, where $\sigma^{(j)} = (1, 2, \dots, n) + \alpha_j$ is the j th adjacent transposition. If C and C^{-1} are the Cartan and inverse Cartan matrices of \mathbf{A}_{n-1} , i.e., $C_{i,j} = 2\delta_{i,j} - \delta_{i,j-1} - \delta_{i,j+1}$ and $C_{i,j}^{-1} = \min\{i, j\} - ij/n$, then $\alpha_i = \sum_{j=1}^{n-1} C_{i,j} \bar{\Lambda}_j$, $(\bar{\Lambda}_i|\alpha_j) = \delta_{i,j}$, $(\alpha_i|\alpha_j) = 2C_{i,j}$ and $(\bar{\Lambda}_i|\bar{\Lambda}_j) = C_{i,j}^{-1}$.

Finally, introducing the notation $(q)\lambda = \prod_{i \geq 1} (q)\lambda_i$ for λ a composition, and using the shorthand $a \equiv b \pmod{c}$ for $a \equiv b \pmod{c}$, our conjecture can be stated as follows.

Conjecture 2.1. *For $\mu \in \mathcal{P}$ and $\nu \in \mathbb{Z}^n$ such that $l(\mu) \leq n$ and $|\mu| = |\nu|$,*

$$K_{\mu\nu} = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \equiv |\mu| \pmod{n}}} \sum_{\sigma \in S_n} \sum_{\lambda \in nQ + \sigma(\rho) - \rho} \epsilon(\sigma) q^{\frac{1}{2n}(\lambda|\lambda + 2\rho)} \\ \times \frac{1}{(q)_{m(\eta)}} \left[\begin{matrix} m(\eta) \\ |\eta| - |\nu| \pmod{n} (1^n) + \nu \end{matrix} \right] \left[\begin{matrix} m(\eta) \\ |\eta| - |\mu| \pmod{n} (1^n) + \mu - \lambda \end{matrix} \right].$$

Since the right side satisfies the periodicity $f_{\mu\nu} = f_{\mu+(a^n), \nu+(b^n)}$ for $a, b \in \mathbb{Z}$, the condition $|\mu| = |\nu|$ cannot be dropped. We also remark that in the notation of [32] $(q)_{m(\eta)} = b_\eta(q)$.

In the following we give proofs of Conjecture 2.1 for $n = 2$ and for $q = 0$.

Proof for $n = 2$. Set $\eta = (1^r)$ (so that $m(\eta) = |\eta| = r$) and $\lambda = (s, -s)$. When $\sigma = (1, 2)$ ($\sigma = (2, 1)$) we need to sum s over the even (odd) integers. Using (2.23) the $n = 2$ case of the conjecture thus becomes

$$K_{\mu\nu} = \sum_{\substack{r=0 \\ r \equiv |\mu| \pmod{2}}}^{\infty} \sum_{s=-\infty}^{\infty} \frac{(-1)^s q^{\binom{s+1}{2}}}{(q)_r} \left[\begin{matrix} r \\ \frac{1}{2}(r + \nu_{12}) \end{matrix} \right]_q \left[\begin{matrix} r \\ \frac{1}{2}(r + \mu_{12}) - s \end{matrix} \right]_q,$$

where $\mu_{12} = \mu_1 - \mu_2$, $\nu_{12} = \nu_1 - \nu_2$ with $\mu = (\mu_1, \mu_2)$ and $\nu = (\nu_1, \nu_2)$ such that $|\mu| = |\nu|$.

Without loss of generality we may assume that $\nu_{12} \geq 0$. Then, by (1.10), the left side is nothing but $\chi(\mu_2 \leq \nu_1 \leq \mu_1) = \chi(\mu_{12} \geq \nu_{12})$. This is also true if $\nu_2 < 0$ as it implies $K_{\mu\nu} = 0$ as well as $\nu_{12} > |\nu| = |\mu| \geq \mu_{12}$.

On the right we now change $s \rightarrow (r + \mu_{12})/2 - s$ and perform the sum over s using the q -binomial theorem [2, Thm. 3.3]

$$(2.27) \quad \sum_{k=0}^n (-z)^k q^{\binom{k}{2}} \left[\begin{matrix} n \\ k \end{matrix} \right]_q = (z)_n.$$

As a result

$$\chi(\mu_{12} \geq \nu_{12}) = \sum_{\substack{r=\nu_{12} \\ r \equiv |\mu| \pmod{2}}}^{\mu_{12}} (-1)^{(r+\mu_{12})/2} q^{\binom{(r+\mu_{12})/2+1}{2}} \frac{(q^{-(r+\mu_{12})/2})_r}{(q)_r} \left[\begin{matrix} r \\ \frac{1}{2}(r + \nu_{12}) \end{matrix} \right]_q.$$

Replacing $r \rightarrow 2r + \nu_{12}$ and making some simplifications yields

$$\begin{aligned} \chi(\mu_{12} \geq \nu_{12}) &= (-1)^{(\mu_{12}-\nu_{12})/2} q^{\binom{\nu_{12}-\mu_{12}}{2}} \left[\begin{matrix} \frac{1}{2}(\mu_{12} + \nu_{12}) \\ \nu_{12} \end{matrix} \right]_q \\ &\quad \times \sum_{r=0}^{(\mu_{12}-\nu_{12})/2} \frac{(q^{(\mu_{12}+\nu_{12})/2+1}, q^{-(\mu_{12}-\nu_{12})/2})_r}{(q, q^{\nu_{12}+1})_r}. \end{aligned}$$

By the q -Chu–Vandermonde sum [13, Eq. (II.7)]

$$(2.28) \quad {}_2\phi_1(a, q^{-n}; c; q, cq^n/a) := \sum_{k=0}^n \frac{(a, q^{-n})_k}{(q, c)_k} \left(\frac{cq^n}{a} \right)^k = \frac{(c/a)_n}{(c)_n}$$

this is readily found to be true. \square

Proof for $q = 0$. Replace $\lambda \rightarrow \sigma(\lambda + \rho) - \rho = \sigma(\lambda + \delta) - \delta$ in the sum on the right. This leaves $(\lambda|\lambda + 2\rho)$ unchanged, yielding

$$(2.29) \quad K_{\mu\nu} = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \equiv |\mu| \pmod{n}}} \sum_{\sigma \in S_n} \sum_{\lambda \in nQ} \epsilon(\sigma) q^{\frac{1}{2n}(\lambda|\lambda+2\rho)} \\ \times \frac{1}{(q)_{m(\eta)}} \left[\begin{matrix} m(\eta) \\ \frac{|\eta|-|\nu|}{n}(1^n) + \nu \end{matrix} \right] \left[\begin{matrix} m(\eta) \\ \frac{|\eta|-|\mu|}{n}(1^n) + \mu - \sigma(\lambda + \delta) + \delta \end{matrix} \right].$$

Now use that (i) $q^{(\lambda|\lambda+2\rho)} = \delta_{\lambda, \emptyset} + O(q)$ for $\lambda \in nQ$, (ii) $\left[\begin{matrix} m(\eta) \\ \mu \end{matrix} \right] = K_{\eta', \mu} + O(q)$, as follows from $K_{\lambda\mu}(0) = \delta_{\lambda, \mu}$ and equation (2.22), and (iii) $1/(q)_{m(\eta)} = 1 + O(q)$. Hence the constant term of (2.29) can be extracted as

$$K_{\mu\nu} = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \equiv |\mu| \pmod{n}}} \sum_{\sigma \in S_n} \epsilon(\sigma) K_{\eta', \frac{|\eta|-|\nu|}{n}(1^n) + \nu} K_{\eta', \frac{|\eta|-|\mu|}{n}(1^n) + \mu - \sigma(\delta) + \delta}.$$

Making the replacement $\sigma \rightarrow \sigma^{-1}$ and using the symmetry (2.5) results in the sum $\sum_{\sigma \in S_n} \epsilon(\sigma) K_{\eta', \sigma(\lambda+\delta) - \delta}$ with $\lambda = \frac{|\eta|-|\mu|}{n}(1^n) + \mu$. By (2.10) (plus subsequent comment) and the substitution $\eta \rightarrow \eta'$ this gives

$$K_{\mu\nu} = \sum_{\substack{\eta \in \mathcal{P} \\ l(\eta) \leq n-1 \\ |\eta| \equiv |\mu| \pmod{n}}} K_{\eta, \frac{|\eta|-|\nu|}{n}(1^n) + \nu} \delta_{\eta, \frac{|\eta|-|\mu|}{n}(1^n) + \mu}.$$

For $\eta, \mu \in \mathcal{P}$ such that $l(\eta), l(\mu) \leq n$, the solution to the linear equation $\eta = \frac{|\eta|-|\mu|}{n}(1^n) + \mu$ is $\eta = \mu + (a^n)$, with a an integer such that $-\mu_n \leq a \leq \eta_n$. Since $\eta_n = 0$, however, a is fixed to $a = -\mu_n$. Hence $\delta_{\eta, \frac{|\eta|-|\mu|}{n}(1^n) + \mu} = \delta_{\eta, \mu - \mu_n(1^n)}$, and after performing the sum over η and using that $|\mu| = |\nu|$ the right-hand side becomes $K_{\mu - \mu_n(1^n), \nu - \mu_n(1^n)}$. Thanks to (2.7) this is $K_{\mu\nu}$. \square

The above proof suggests that what is needed to prove Conjecture 2.1 is a generalization of (2.10) for the sum

$$(2.30) \quad \sum_{\sigma \in S_n} \epsilon(\sigma) K_{\lambda, \sigma(\mu+\delta) - \delta - \alpha}$$

with $\alpha \in Q$. Indeed, substituting (2.22) into (2.29) we obtain

$$K_{\mu\nu} = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \equiv |\mu| \pmod{n}}} \sum_{\omega \vdash |\eta|} \frac{K_{\omega, \eta}(q)}{(q)_{m(\eta)}} \left[\begin{matrix} m(\eta) \\ \frac{|\eta| - |\mu|}{n} (1^n) + \nu \end{matrix} \right] \\ \times \sum_{\lambda \in nQ} q^{\frac{1}{2n}(\lambda|\lambda + 2\rho)} \sum_{\sigma \in S_n} \epsilon(\sigma) K_{\omega, \frac{|\eta| - |\mu|}{n} (1^n) + \mu - \sigma(\lambda + \delta) + \delta}.$$

Changing $\sigma \rightarrow \sigma^{-1}$ and using (2.5) the sum over σ takes exactly the form of (2.30) (with $\mu \rightarrow \frac{|\eta| - |\mu|}{n} (1^n) + \mu$ and $\alpha \rightarrow \lambda$).

We also remark that the proof of the constant term cannot (easily) be extended to deal with low-order terms in q . This because $(\lambda|\lambda + 2\rho) = 2n$ for $\lambda = -n(\bar{\Lambda}_1 + \bar{\Lambda}_{n-1})$, so that restricting the sum over λ to the single term $\lambda = \emptyset$ is only correct to zeroth order.

Assuming Conjecture 2.1 it is not hard to prove the following orthogonality relation.

Corollary 2.2. *For $\mu, \nu \in \mathcal{P}$ such that $l(\mu), l(\nu) \leq n$ and $|\mu| = |\nu|$,*

$$\sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \equiv |\mu| \pmod{n}}} \sum_{\sigma \in S_n} \sum_{\lambda \in nQ + \sigma(\rho) - \rho} \epsilon(\sigma) q^{\frac{1}{2n}(\lambda|\lambda + 2\rho)} \\ \times \frac{K_{\left(\frac{|\eta| - |\nu|}{n} (1^n) + \nu\right)', \eta}(q)}{(q)_{m(\eta)}} \left[\begin{matrix} m(\eta) \\ \frac{|\eta| - |\mu|}{n} (1^n) + \mu - \lambda \end{matrix} \right] = \delta_{\mu, \nu}.$$

Since $K_{\lambda\mu}(q) = 0$ if $\lambda \notin \mathcal{P}$ one may add the additional restriction $|\eta| \geq |\nu| - n\nu_n = \sum_{i=1}^{n-1} (\nu_i - \nu_{i+1})$ to the sum over η . We also note that care should be taken when writing $\left(\frac{|\eta| - |\nu|}{n} (1^n) + \nu\right)'$ as $(n^{(|\eta| - |\nu|)/n}) \cup \nu'$ since $|\eta| - |\nu|$ need not be nonnegative.

Proof. Replace $\nu \rightarrow \tau(\hat{\nu} + \delta) - \delta$ in Conjecture 2.1 with $\hat{\nu} \in \mathcal{P}$. Then multiply the result by $\epsilon(\tau)$ and sum τ over S_n . By (2.10) and (2.26) Corollary 2.2 (with ν replaced by $\hat{\nu}$) follows. \square

A proof very similar to the proof of the $n = 2$ case of Conjecture 2.1 shows that Corollary 2.2 may be viewed as an A_{n-1} generalization of the well-known ${}_2\phi_1(a, q^{-n}; aq^{1-n}; q, q) = \delta_{n,0}$, which corresponds to the specialization $c = aq^{1-n}$ in the q -Chu–Vandermonde sum (2.28). Indeed, if we take $n = 2$ in Corollary 2.2 and sum over σ and λ by the q -binomial theorem (2.27), then the remaining sum over η takes the form of the above ${}_2\phi_1$ sum with $n = \mu_1 - \nu_1$ and $a = q^{\mu_1 - \nu_2 + 1}$.

3. AN A_{n-1} BAILEY LEMMA

3.1. Definitions and main result. Let $\ell \in \mathbb{Z}_+$, $k \in Q \cap P_+$ (i.e., $k = (k_1, \dots, k_n) \in \mathbb{Z}_+^n$ such that $k_1 \geq k_2 \geq \dots \geq k_n$, and $|k| = 0$) and $\eta \in \mathcal{P}$ such that $\eta_1 \leq n - 1$ and $|\eta| \equiv \ell \pmod{n}$. Then $\alpha = (\alpha_k)$ and $\beta = (\beta_\eta)$ form an A_{n-1} Bailey pair relative to q^ℓ if

$$(3.1) \quad \beta_\eta = \sum_{\substack{k \in Q \cap P_+ \\ k_1 \leq (|\eta| - \ell)/n}} \frac{\alpha_k}{(q)_{m(\eta)}} \left[\begin{matrix} m(\eta) \\ \frac{|\eta|}{n} (1^n) - k - \ell \bar{\Lambda}_{n-1} \end{matrix} \right],$$

and $\gamma = (\gamma_k)$ and $\delta = (\delta_\eta)$ form an A_{n-1} conjugate Bailey pair relative to q^ℓ if

$$(3.2) \quad \gamma_k = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \geq \ell + nk_1 \pmod{n}}} \frac{\delta_\eta}{(q)_{m(\eta)}} \left[\begin{matrix} m(\eta) \\ \frac{|\eta|}{n} (1^n) - k - \ell \bar{\Lambda}_{n-1} \end{matrix} \right].$$

Here, and elsewhere in the paper, $a \geq b \pmod{n}$ stands for $a \equiv b \pmod{n}$ such that $a \geq b$. Note that a necessary condition for the above summands to be non-vanishing is $\frac{|\eta|}{n}(1^n) - k - \ell \bar{\Lambda}_{n-1} \in \mathbb{Z}_+^n$. Since $k \in Q \cap P_+$ this boils down to the single condition $(|\eta| - \ell)/n - k_1 \geq 0$, justifying the restrictions in the sums over k and η .

Recalling (2.22), the definition (3.1) is a multivariable generalization of (1.12) up to an irrelevant normalization factor $(q)_\ell$. Of course, to really reduce to (1.12) for $n = 2$ we have to write $\eta = (1^{2L+\ell})$ and then replace $\beta_{(1^{2L+\ell})}$ by β_L . In exactly the same way (3.2) is a (normalized) multivariable generalization of (1.2).

Given an A_{n-1} Bailey pair and conjugate Bailey pair relative to q^ℓ , it follows that

$$(3.3) \quad \sum_{k \in Q \cap P_+} \alpha_k \gamma_k = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \geq \ell \pmod{n}}} \beta_\eta \delta_\eta.$$

If for $n = 2$ we set $\eta = (1^{2L+\ell})$ and write β_L and δ_L instead of β_η and δ_η the right side simplifies to $\sum_{L \geq 0} \beta_L \delta_L$ in accordance with (1.3).

The main result of this section will be an A_{n-1} , level- N generalization of Bailey's conjugate pair (1.4), resulting in an according generalization of Bailey's identity (1.5). First, however, a few more definitions are required.

Let N be a fixed, nonnegative integer, which will be referred to as the level. The canonical basis vectors in \mathbb{R}^{n-1} and \mathbb{R}^{N-1} will be denoted by e_a and \bar{e}_j , respectively, and we set $\bar{e}_0 = \bar{e}_N = \emptyset$. The $(n-1)$ - and $(N-1)$ -dimensional identity matrices are denoted by I and \bar{I} and the A_{N-1} Cartan matrix by \bar{C} . Hence $\bar{C}_{j,k}^{-1} = \min\{j, k\} - jk/N$, which will also be used when either j or k is 0 or N . For $u, v \in \mathbb{Z}^k$ with k either $n-1$, $N-1$ or $(n-1)(N-1)$ and M an k -dimensional (square) matrix, Mu is the vector with components $(Mu)_i = \sum_{j=1}^k M_{i,j} u_j$ and $vMu = \sum_{i,j=1}^k v_i M_{i,j} u_j$. We already defined $(q)_u = \prod_{i=1}^k (q)_{u_i}$ and extending this, we also use $\begin{bmatrix} u+v \\ u \end{bmatrix}_q = \prod_{i=1}^k \begin{bmatrix} u_i+k_i \\ u_i \end{bmatrix}_q$. Given integers $m_j^{(a)}$ for $a \in [n-1]$ and $j \in [N-1]$, the vector m is defined as

$$(3.4) \quad m = \sum_{a=1}^{n-1} \sum_{j=1}^{N-1} (e_a \otimes \bar{e}_j) m_j^{(a)} \in \mathbb{Z}^{(n-1)(N-1)}.$$

Let $\eta, \nu \in \mathcal{P}$ such that $\eta_1 \leq n-1$, $\nu_1 \leq N$ and $|\eta| \equiv |\nu| \pmod{n}$, and set $\mu = \sum_{a=1}^{n-1} m_a(\eta) \bar{\Lambda}_a = \sum_{a=1}^{n-1} (\eta'_a - \eta'_{a+1}) \bar{\Lambda}_a \in P_+$ ($\eta'_n = 0$), so that $|\nu| \bar{\Lambda}_1 - \mu \in Q$. Then the polynomial $F_{\eta\nu}(q)$ is defined by

$$(3.5) \quad F_{\eta\nu}(q) = q^{n(\nu) - \frac{(n-1)|\nu|^2}{2nN} + \frac{(\mu|\mu)}{2N}} \sum_m q^{\frac{1}{2}m(C \otimes \bar{C}^{-1})m - \sum_{i=1}^{l(\nu)} (e_1 \otimes \bar{e}_{N-\nu_i})(I \otimes \bar{C}^{-1})m} \begin{bmatrix} m+p \\ m \end{bmatrix}_q.$$

Here the vector $p \in \mathbb{Z}^{(n-1)(N-1)}$ is determined by

$$(3.6) \quad (C \otimes \bar{I})m + (I \otimes \bar{C})p = (m(\eta) \otimes \bar{e}_1) + \sum_{i=1}^{l(\nu)} (e_1 \otimes \bar{e}_{N-\nu_i})$$

and the sum is over m such that

$$(3.7) \quad \sum_{a=1}^{n-1} \sum_{j=1}^{N-1} j m_j^{(a)} \alpha_a \in NQ + \mu - |\nu| \bar{\Lambda}_1.$$

This last equation ensures that the components of p are integers, but the reverse of this is not true; demanding that $p_j^{(a)} \in \mathbb{Z}$ leads to a restriction on m that, generally, is weaker than (3.7). We also caution the reader not to confuse $m(\eta) \in \mathbb{Z}_+^{n-1}$ with the vector $m \in \mathbb{Z}^{(n-1)(N-1)}$.

For our next definition it will be convenient to view μ rather than η as primary variable, and for $\mu \in P_+$ and $\nu \in \mathcal{P}$ such that $\nu_1 \leq N$ and $|\nu| \bar{\Lambda}_1 - \mu \in Q$ we define

$$(3.8) \quad C_{\mu\nu}(q) = \frac{q^{n(\nu) - \frac{(n-1)|\nu|^2}{2nN} + \frac{(\mu|\mu)}{2N}}}{(q)_\infty^{n-1}} \sum_m \frac{q^{\frac{1}{2}m(C \otimes \bar{C}^{-1})m - \sum_{i=1}^{l(\nu)} (e_1 \otimes \bar{e}_{N-\nu_i})(I \otimes \bar{C}^{-1})m}}{(q)_m},$$

where the sum is again over m such that (3.7) holds.

From (2.1) it readily follows that both $F_{\eta\nu}$ and $C_{\mu\nu}$ satisfy the equation $g_{\xi, (N^n) \cup \nu}(q) = q^{|\nu|} g_{\xi\nu}(q)$. Without loss of generality we may therefore assume that ν has at most $n-1$ parts equal to N . From (3.4) it also follows that $F_{\eta\nu}$ and $C_{\mu\nu}$ trivialize for $N=1$ to

$$(3.9) \quad F_{\eta, (1^i)}(q) = q^{\frac{1}{2}(\mu|\mu) - \frac{1}{2}\|\bar{\Lambda}_i\|^2}$$

with $\mu = \sum_{a=1}^{n-1} m_a(\eta) \bar{\Lambda}_a$ and

$$(3.10) \quad C_{\mu, (1^i)}(q) = \frac{q^{\frac{1}{2}(\mu|\mu) - \frac{1}{2}\|\bar{\Lambda}_i\|^2}}{(q)_\infty^{n-1}},$$

where $i \in \{0, \dots, n-1\}$. For later use it will also be convenient to define

$$(3.11) \quad F_{\eta, \emptyset}(q) = \delta_{\eta, \emptyset} \quad \text{and} \quad C_{\mu, \emptyset}(q) = \delta_{\mu, \emptyset} \quad \text{for } N=0.$$

In the important special case $\nu = (sN^r)$, the polynomial $F_{\eta\nu}(q)$ can be identified with a level- N restricted version of the generalized Kostka polynomials $K_{\lambda R}(q)$ [41]. Denoting these polynomials by $K_{\lambda R}^N(q)$ it follows from [41, Eq. (6.7)] that for $\nu = (sN^r)$ with $s \in \{0, \dots, N\}$

$$F_{\eta\nu}(q) = q^{\sum_{a=1}^{n-1} \binom{\eta'_a}{2} + s + \frac{n-r}{n}(|\eta| - |\nu|)} K_{\lambda R}^N(q^{-1}),$$

where $\lambda = \frac{1}{n}(nN + |\eta| - |\nu|)(1^n) + (s)$ and R is a sequence of $|m(\eta)| + n - r$ partitions with $m_a(\eta)$ partitions (1^a) ($a \in [n-1]$) and $n-r$ partitions (N) .

Finally we state our main result, which for $n \geq 3$ was conjectured in [44, Eq. (9.9)] (see also [52, Conj. 14]). For $n=2$ the theorem below is equivalent to the ‘‘higher-level Bailey lemma’’ of [42, Cor. 4.1], and for $N=1$ and $\ell=0$ (but general n) it was proven in [52, Prop. 12].

Theorem 3.1. *Let $\ell \in \mathbb{Z}_+$, $k \in Q \cap P_+$ and $\eta, \nu \in \mathcal{P}$ such that $\nu_1 \leq N$, $\eta_1 \leq n-1$ and $|\eta| \equiv |\nu| \equiv \ell \pmod{n}$. Then*

$$\gamma_k = C_{\ell \bar{\Lambda}_1 - \pi(k), \nu}(q) \quad \text{and} \quad \delta_\eta = F_{\eta\mu}(q)$$

form an A_{n-1} conjugate Bailey pair relative to q^ℓ .

Substituting the conjugate Bailey pair of Theorem 3.1 into (3.3) we obtain the following generalization of Bailey’s key-identity (1.5):

$$(3.12) \quad \sum_{k \in Q \cap P_+} \alpha_k C_{\ell \bar{\Lambda}_1 - \pi(k), \nu}(q) = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \geq \ell \pmod{n}}} \beta_\eta F_{\eta\nu}(q).$$

Equation (1.5) with $a = q^\ell$ follows for $n=2$ and $N=1$. According to (3.9) and (3.10) one then finds $C_{(\ell/2+L, \ell/2-L), (1^i)}(q) = g_{i, \ell}(q) q^{L(L+\ell)} / (q)_\infty$ and $F_{(1^{2L+\ell}), (1^i)}(q) = g_{i, \ell}(q) q^{L(L+\ell)}$, where $g_{i, \ell}(q) = q^{i(i-2)/4 + \ell^2/4}$ is an irrelevant factor that drops out of (3.12).

Applications of (3.12) will be given in Section 6, with the remainder of this section devoted to a proof of Theorem 3.1. This proof consists of several steps. First, in Lemma 3.2 below, we reformulate the definition of an A_{n-1} conjugate Bailey pair. Then we state Theorem 3.3, which claims a deep identity for Kostka polynomials. A straightforward specialization leads to (3.13), which in combination with Lemma 3.2 proves Theorem 3.1. To prove Theorem 3.3 we need the Propositions 3.4 and 3.5 below, which are due to Hatayama *et al.* [15] (see also [16]).

Because of their importance to Theorem 3.3 we have included Sections 3.3 and 3.4 containing proofs of both propositions.

3.2. Proof of Theorem 3.1.

Lemma 3.2. *The definition (3.2) of an A_{n-1} conjugate Bailey pair (relative to q^ℓ) can be rewritten as*

$$\gamma_k = \sum_{\substack{\lambda \in \mathcal{P} \\ l(\lambda) \leq n-1 \\ |\lambda| \geq \ell + nk_1}} K_{\lambda, \frac{|\lambda|}{n}(1^n) - k - \ell \bar{\Lambda}_{n-1}} \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \geq |\lambda|}} \frac{\delta_\eta K_{(n(|\eta| - |\lambda|)/n) \cup \lambda', \eta}(q)}{(q)_{m(\eta)}}.$$

Proof. Using (2.22) to eliminate the antisymmetric supernomial from (3.2) yields

$$\gamma_k = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \geq \ell + nk_1}} \frac{\delta_\eta}{(q)_{m(\eta)}} \sum_{\nu \vdash |\eta|} K_{\nu, \frac{|\nu|}{n}(1^n) - k - \ell \bar{\Lambda}_{n-1}} K_{\nu', \eta}(q).$$

By (2.5), (2.6) and $\min(|\eta|(1^n)/n - k - \ell \bar{\Lambda}_{n-1}) = (|\eta| - \ell)/n - k_1$ (since $k \in Q \cap P_+$) we may add the conditions $l(\nu) \leq n$ and $(|\eta| - \ell)/n - k_1 - \nu_n \in \mathbb{Z}_+$ to the sum over ν . Changing the order of summation then gives

$$\gamma_k = \sum_{\substack{\nu \in \mathcal{P} \\ l(\nu) \leq n \\ |\nu| - n\nu_n \geq \ell + nk_1}} K_{\nu, \frac{|\nu|}{n}(1^n) - k - \ell \bar{\Lambda}_{n-1}} \sum_{\substack{\eta \vdash |\nu| \\ \eta_1 \leq n-1}} \frac{\delta_\eta K_{\nu', \eta}(q)}{(q)_{m(\eta)}}.$$

Writing $\nu = (a^n) + \lambda$ with λ a partition such that $l(\lambda) \leq n-1$ this becomes

$$\begin{aligned} \gamma_k &= \sum_{a=0}^{\infty} \sum_{\substack{\lambda \in \mathcal{P} \\ l(\lambda) \leq n-1 \\ |\lambda| \geq \ell + nk_1}} K_{(a^n) + \lambda, (a^n) + \frac{|\lambda|}{n}(1^n) - k - \ell \bar{\Lambda}_{n-1}} \sum_{\substack{\eta \vdash a^n + |\lambda| \\ \eta_1 \leq n-1}} \frac{\delta_\eta K_{(n^a) \cup \lambda', \eta}(q)}{(q)_{m(\eta)}} \\ &= \sum_{\substack{\lambda \in \mathcal{P} \\ l(\lambda) \leq n-1 \\ |\lambda| \geq \ell + nk_1}} K_{\lambda, \frac{|\lambda|}{n}(1^n) - k - \ell \bar{\Lambda}_{n-1}} \sum_{a=0}^{\infty} \sum_{\substack{\eta \vdash a^n + |\lambda| \\ \eta_1 \leq n-1}} \frac{\delta_\eta K_{(n^a) \cup \lambda', \eta}(q)}{(q)_{m(\eta)}}, \end{aligned}$$

where the second equality follows from (2.7). Interchanging the sums over η and a it follows that a is fixed by $a = (|\eta| - |\lambda|)/n$, resulting in the claim of the lemma. \square

Theorem 3.3. *For $\mu \in P_+$ and $\nu \in \mathcal{P}$ such that $\nu_1 \leq N$ and $|\nu| \bar{\Lambda}_1 - \mu \in Q$,*

$$\sum_{\substack{\lambda \in \mathcal{P} \\ l(\lambda) \leq n-1 \\ |\lambda| \bar{\Lambda}_1 - \mu \in Q}} K_{\lambda, \frac{|\lambda|}{n}(1^n) + \mu} \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \geq |\lambda|}} \frac{F_{\eta\nu}(q) K_{(n(|\eta| - |\lambda|)/n) \cup \lambda', \eta}(q)}{(q)_{m(\eta)}} = C_{\mu\nu}(q).$$

Before proving this, let us show how combined with Lemma 3.2 it implies Theorem 3.1. By (2.5), μ on the left may be replaced by $\pi(\mu)$. After this change we choose $\mu = \ell \bar{\Lambda}_1 - \pi(k)$ with $k \in Q \cap P_+$ and $\ell \in \mathbb{Z}_+$, and use that $\pi(\bar{\Lambda}_1) = -\bar{\Lambda}_{n-1}$ and $(\bar{\Lambda}_1)_a - (\bar{\Lambda}_1)_{a+1} = \delta_{a,1}$. Hence

$$\begin{aligned} (3.13) \quad & \sum_{\substack{\lambda \in \mathcal{P} \\ l(\lambda) \leq n-1 \\ |\lambda| \geq \ell + nk_1}} K_{\lambda, \frac{|\lambda|}{n}(1^n) - k - \ell \bar{\Lambda}_{n-1}} \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \geq |\lambda|}} \frac{F_{\eta\nu}(q) K_{(n(|\eta| - |\lambda|)/n) \cup \lambda', \eta}(q)}{(q)_{m(\eta)}} \\ & = C_{\ell \bar{\Lambda}_1 - \pi(k), \nu}(q) \end{aligned}$$

with $|\nu| \equiv \ell \pmod{n}$. Comparing this with Lemma 3.2 yields Theorem 3.1.

The proof of Theorem 3.3 requires the following two propositions.

Proposition 3.4. *For $\lambda, \nu \in \mathcal{P}$ such that $l(\lambda) \leq n$, $\nu_1 \leq N$ and $|\lambda| = |\nu|$,*

$$\lim_{M \rightarrow \infty} q^{-nN \binom{M}{2} - M|\nu|} K_{NM(1^n) + \lambda, (N^{nM}) \cup \nu}(q) = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \equiv |\lambda| \pmod{n}}} \frac{F_{\eta\nu}(q) K_{\xi'\eta}(q)}{(q)_{m(\eta)}},$$

where $\xi = \frac{|\eta| - |\lambda|}{n}(1^n) + \lambda$.

Proposition 3.5. *For $\lambda, \nu \in \mathcal{P}$ such that $l(\lambda) \leq n$, $\nu_1 \leq N$ and $|\lambda| = |\nu|$,*

$$\lim_{M \rightarrow \infty} q^{-nN \binom{M}{2} - M|\nu|} \mathcal{S}_{NM(1^n) + \lambda, (N^{nM}) \cup \nu}(q) = C_{\mu\nu}(q),$$

where $\mu = \sum_{a=1}^{n-1} (\lambda_a - \lambda_{a+1}) \bar{\Lambda}_a \in P_+$.

Proposition 3.4 is [15, Prop. 6.4] and Proposition 3.5 follows from [15, Props. 4.3, 4.6 and 5.8]. Before giving proofs we first show how these results imply Theorem 3.3.

Proof of Theorem 3.3. For $\mu \in P$ and $\nu \in \mathcal{P}$ such that $\nu_1 \leq N$ and $|\nu| \bar{\Lambda}_1 - \mu \in Q$, consider the expression

$$(3.14) \quad I_{\mu\nu}(q) = \lim_{M \rightarrow \infty} q^{-nN \binom{M}{2} - M|\nu|} \sum_{\substack{\lambda \in \mathcal{P} \\ l(\lambda) \leq n-1 \\ |\lambda| \bar{\Lambda}_1 - \mu \in Q}} K_{\lambda, \frac{|\lambda|}{n}(1^n) + \mu} K_{\frac{nNM + |\nu| - |\lambda|}{n}(1^n) + \lambda, (N^{nM}) \cup \nu}(q).$$

By Proposition 3.4 with $\lambda \rightarrow \frac{1}{n}(|\nu| - |\lambda|)(1^n) + \lambda$ it is readily found that

$$I_{\mu\nu}(q) = \sum_{\substack{\lambda \in \mathcal{P} \\ l(\lambda) \leq n-1 \\ |\lambda| \bar{\Lambda}_1 - \mu \in Q}} K_{\lambda, \frac{|\lambda|}{n}(1^n) + \mu} \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \geq |\lambda| \pmod{n}}} \frac{F_{\eta\nu}(q) K_{(n(|\eta| - |\lambda|)/n) \cup \lambda', \eta}(q)}{(q)_{m(\eta)}}.$$

Here we have used that $\xi = \frac{|\eta| - |\lambda|}{n}(1^n) + \lambda$ with $l(\lambda) \leq n-1$ is a partition iff $|\eta| \geq |\lambda| \pmod{n}$. Hence we may assume this condition and write $\xi' = (n(|\eta| - |\lambda|)/n) \cup \lambda'$.

In order to prove Theorem 3.3 it remains to be shown that $I_{\mu\nu} = C_{\mu\nu}$ if $\mu \in P_+$. To this end we use the properties (2.7) and (2.6) of the Kostka numbers to identify the summand of (3.14) with a completely symmetric supernomial;

$$\begin{aligned} & \sum_{\substack{\lambda \in \mathcal{P} \\ l(\lambda) \leq n-1 \\ |\lambda| \bar{\Lambda}_1 - \mu \in Q}} K_{\lambda, \frac{|\lambda|}{n}(1^n) + \mu} K_{\frac{nNM + |\nu| - |\lambda|}{n}(1^n) + \lambda, (N^{nM}) \cup \nu}(q) \\ &= \sum_{\substack{\lambda \in \mathcal{P} \\ l(\lambda) \leq n-1 \\ |\lambda| \bar{\Lambda}_1 - \mu \in Q}} K_{\frac{nNM + |\nu| - |\lambda|}{n}(1^n) + \lambda, \frac{nNM + |\nu|}{n}(1^n) + \mu} \\ & \quad \times K_{\frac{nNM + |\nu| - |\lambda|}{n}(1^n) + \lambda, (N^{nM}) \cup \nu}(q) \\ &= \sum_{\substack{\eta \vdash |\nu| + nNM \\ l(\eta) \leq n}} K_{\eta, \frac{nNM + |\nu|}{n}(1^n) + \mu} K_{\eta, (N^{nM}) \cup \nu}(q) \\ &= \mathcal{S}_{\frac{nNM + |\nu|}{n}(1^n) + \mu, (N^{nM}) \cup \nu}(q). \end{aligned}$$

The last equality follows from (2.21) and the fact that the restriction $l(\lambda) \leq n$ in the second-last line may be dropped thanks to (2.5) and (2.6).

By Proposition 3.5 with $\lambda \rightarrow \frac{|\nu|}{n}(1^n) + \mu$ (which is a partition iff $\mu \in P_+$) it thus follows that for $\mu \in P_+$ there holds $I_{\mu\nu} = C_{\mu\nu}$. \square

3.3. Proof of Proposition 3.4. As mentioned previously, Proposition 3.4 is due to Hatayama *et al.* [15, Prop. 6.4], who we will closely follow in our proof. The only significant difference is that in [15] only the case $\nu = \emptyset$ is treated in detail.

Key to Proposition 3.4 are two so-called fermionic representations of the Kostka polynomials. The first of these is due to Kirillov and Reshetikhin [24, Thm. 4.2]. Let $T_{j,k} = \min\{j, k\}$.

Proposition 3.6. *For $\lambda, \mu \in \mathcal{P}$ such that $|\lambda| = |\mu|$ and $l(\lambda) \leq n$,*

$$(3.15) \quad K_{\lambda\mu}(q) = q^{n(\mu)} \sum q^{\frac{1}{2} \sum_{a,b=1}^{n-1} \sum_{j,k \geq 1} m_j^{(a)} C_{a,b} T_{j,k} m_k^{(b)}} \\ \times q^{-\sum_{i=1}^{l(\mu)} \sum_{j \geq 1} T_{\mu_i, j} m_j^{(1)}} \prod_{a=1}^{n-1} \prod_{j \geq 1} \left[\begin{matrix} m_j^{(a)} + p_j^{(a)} \\ m_j^{(a)} \end{matrix} \right]_q.$$

Here $p_j^{(a)}$ is given by

$$(3.16) \quad p_j^{(a)} = \delta_{a,1} \sum_{i=1}^{l(\mu)} T_{j, \mu_i} - \sum_{b=1}^{n-1} \sum_{k \geq 1} C_{a,b} T_{j,k} m_k^{(b)}$$

and the sum is over $m_j^{(a)} \in \mathbb{Z}_+$ for $a \in [n-1]$ and $j \geq 1$, subject to

$$(3.17) \quad \sum_{j \geq 1} j m_j^{(a)} = \sum_{b=a+1}^n \lambda_b.$$

Using (i) the duality (2.18), (ii) the analogue of Proposition 3.6 for generalized Kostka polynomials, conjectured in [26, Conj. 6] and [44, Conj. 8.3] and proven in [25, Thm 2.10], and (iii) (2.14), a second fermionic representation of the Kostka polynomials arises [15, Prop. 5.6].

Proposition 3.7. *For $\lambda, \eta \in \mathcal{P}$ such that $|\lambda| = |\eta|$, $l(\lambda) \leq n$ and $\eta_1 \leq n-1$*

$$(3.18) \quad K_{\lambda'\eta}(q) = \sum q^{\frac{1}{2} \sum_{a,b=1}^{n-1} \sum_{j,k \geq 1} m_j^{(a)} C_{a,b} T_{j,k} m_k^{(b)}} \prod_{a=1}^{n-1} \prod_{j \geq 1} \left[\begin{matrix} m_j^{(a)} + p_j^{(a)} \\ m_j^{(a)} \end{matrix} \right]_q.$$

Here $p_j^{(a)}$ is given by

$$(3.19) \quad p_j^{(a)} = m_a(\eta) - \sum_{b=1}^{n-1} \sum_{k \geq 1} C_{a,b} T_{j,k} m_k^{(b)},$$

the sum is over $m_j^{(a)} \in \mathbb{Z}_+$ for $a \in [n-1]$ and $j \geq 1$ such that

$$(3.20) \quad \sum_{j \geq 1} j m_j^{(a)} = (\bar{\Lambda}_a | \mu) - \sum_{b=1}^a \left(\lambda_b - \frac{|\lambda|}{n} \right)$$

and $\mu = \sum_{a=1}^{n-1} m_a(\eta) \bar{\Lambda}_a$.

As an example, let us calculate $K_{(4,3,1), (3,3,2)}(q)$. Taking the Kirillov–Reshetikhin representation we have to compute all solutions to $\sum_{j \geq 1} j m_j^{(1)} = 4$ and $\sum_{j \geq 1} m_j^{(2)} = 1$. The first of these equations has five solutions corresponding to the five partitions of 4. The second equation has the unique solution $m_j^{(2)} = \delta_{j,1}$. Calculating the corresponding $p_j^{(a)}$ for each of the five solutions using (3.16), it turns out that only one of the five has all $p_j^{(a)}$ nonnegative. Hence

only this solution, given by $m_1^{(1)} = m_3^{(1)} = m_1^{(2)} = 1$, $m_j^{(a)} = 0$ otherwise and $p_j^{(1)} = \chi(j \geq 2)$, $p_j^{(2)} = p_j^{(1)} + \chi(j \geq 3)$ contributes to the sum yielding $K_{(4,3,1),(3,3,2)}(q) = q \begin{bmatrix} 2 \\ 1 \end{bmatrix}_q = q + q^2$.

Taking the representation of Proposition 3.7 we have $\lambda' = (4, 3, 1)$, $\eta = (3, 3, 2)$ and hence $\lambda = (3, 2, 2, 1)$, $m(\eta) = (0, 1, 2)$. We may thus take $n = 4$ yielding $\mu = (1, 1, 0, -2)$ and leading to the equations $\sum_{j \geq 1} j m_j^{(1)} = 0$ and $\sum_{j \geq 1} j m_j^{(a)} = 1$ for $a \in \{2, 3\}$. This has the unique solution $m_1^{(2)} = m_1^{(3)} = 1$ and $m_j^{(a)} = 0$ otherwise. From (3.19) it then follows that $p_j^{(1)} = p_j^{(3)} = 1$ and $p_j^{(2)} = 0$ for all $j \geq 1$, so that, once more, $K_{(4,3,1),(3,3,2)}(q) = q \begin{bmatrix} 2 \\ 1 \end{bmatrix}_q = q + q^2$.

To now prove Proposition 3.4, take the Kostka polynomial in the representation of Proposition 3.6, assume that $\mu_1 \leq N$, and replace

$$(3.21) \quad \lambda \rightarrow NM(1^n) + \lambda \quad \text{and} \quad \mu \rightarrow (N^{nM}) \cup \nu$$

with $\lambda, \nu \in \mathcal{P}$ such that $l(\lambda) \leq n$, $\nu_1 \leq N$ and $|\lambda| = |\nu|$. By this change equations (3.16) and (3.17) become

$$(3.22) \quad p_j^{(a)} = \delta_{a,1} \sum_{i=1}^{l(\nu)} T_{j,\nu_i} - \sum_{b=1}^{n-1} \sum_{k \geq 1} C_{a,b} T_{j,k} (m_k^{(b)} - v_k^{(b)})$$

and

$$(3.23) \quad \sum_{j \geq 1} j (m_j^{(a)} - v_j^{(a)}) = \sum_{b=a+1}^n \lambda_b,$$

where $v_j^{(a)} = M(n-a)\delta_{j,N}$. Without loss of generality we may assume that $p_N^{(a)} \geq 0$, and we define $\eta \in \mathcal{P}$ such that $\eta_1 \leq n-1$ by $m_a(\eta) = p_N^{(a)}$ and set $\mu = \sum_{a=1}^{n-1} m_a(\eta) \bar{\Lambda}_a \in P_+$. Then (3.22) implies

$$(3.24) \quad m_a(\eta) = |\nu| \delta_{a,1} - \sum_{b=1}^{n-1} \sum_{j \geq 1} C_{a,b} T_{N,j} (m_j^{(b)} - v_j^{(b)}).$$

By $|\eta| = \sum_{a=1}^{n-1} a m_a(\eta)$ and $\sum_{a=1}^{n-1} a C_{a,b} = n \delta_{b,n-1}$ it follows that $\eta \equiv |\nu| \pmod{n}$. The equations (3.22), (3.24) and $T_{j,k} = \chi(k \in [N-1]) \bar{C}_{j,k}^{-1} + j T_{N,k}/N$ (true for $k \geq 0$ and $0 \leq j \leq N$) further yield

$$(3.25) \quad \begin{aligned} p_j^{(a)} &= \delta_{a,1} \sum_{i=1}^{l(\nu)} T_{j,\nu_i} - \sum_{b=1}^{n-1} C_{a,b} \left[\sum_{k=1}^{N-1} \bar{C}_{j,k}^{-1} m_k^{(b)} + \frac{j}{N} \sum_{k \geq 1} T_{N,k}^{-1} (m_k^{(b)} - v_k^{(b)}) \right] \\ &= \delta_{a,1} \sum_{i=1}^{l(\nu)} \bar{C}_{j,\nu_i}^{-1} + \frac{j}{N} m_a(\eta) - \sum_{b=1}^{n-1} \sum_{k=1}^{N-1} C_{a,b} \bar{C}_{j,k}^{-1} m_k^{(b)} \end{aligned}$$

for $j \in [N-1]$. Similarly, by (3.22), $\nu_1 \leq N$ and $T_{j+N,k} = \chi(k > N) T_{j,k-N} + T_{k,N}$ (true for $j \geq 0$) one finds

$$(3.26) \quad \begin{aligned} p_{j+N}^{(a)} &= |\nu| \delta_{a,1} - \sum_{b=1}^{n-1} C_{a,b} \left[\sum_{k \geq 1} T_{j,k} m_{k+N}^{(b)} + \sum_{k \geq 1} T_{k,N} (m_k^{(b)} - v_k^{(b)}) \right] \\ &= m_a(\eta) - \sum_{b=1}^{n-1} \sum_{k \geq 1} C_{a,b} T_{j,k} m_{k+N}^{(b)} \end{aligned}$$

for $j \geq 1$. Note that (3.25) for $j = N$ and (3.26) for $j = 0$ correspond to the tautology $p_N^{(a)} = m_a(\eta)$.

Equation (3.24), the definition of μ and $|\nu| = |\lambda|$ may also be applied to rewrite the restriction (3.23). Namely, if in (3.24) we replace $a \rightarrow d$, then multiply by $C_{a,d}^{-1}$ and sum over d , and finally subtract (3.23) from the resulting equation, we find

$$(3.27) \quad \sum_{j \geq 1} j m_{j+N}^{(a)} = (\bar{\Lambda}_a |\mu|) - \sum_{b=1}^a \left(\lambda_b - \frac{|\lambda|}{n} \right).$$

Finally consider the exponent E of q on the right of (3.15) after the replacements (3.21). From (3.24) and (2.1) it follows that

$$\begin{aligned} E &= nN \binom{M}{2} + M|\nu| + n(\nu) - \frac{(n-1)|\nu|^2}{2nN} + \frac{(\mu|\mu)}{2N} \\ &\quad + \frac{1}{2} \sum_{a,b=1}^{n-1} \sum_{j,k=1}^{N-1} m_j^{(a)} C_{a,b} \bar{C}_{j,k}^{-1} \left(m_k^{(b)} - \sum_{i=1}^{l(\nu)} \delta_{a,1} \delta_{b,1} \delta_{k,\nu_i} \right) \\ &\quad + \frac{1}{2} \sum_{a,b=1}^{n-1} \sum_{j,k \geq 1} C_{a,b} \bar{C}_{j,k}^{-1} m_{j+N}^{(a)} m_{k+N}^{(b)}. \end{aligned}$$

This result does not contain $m_N^{(a)}$, and in order to take the large M limit we replace $m_j^{(a)} \rightarrow m_j^{(a)} + v_j^{(a)} = m_j^{(a)} + M(n-a)\delta_{j,N}$. Then the M -dependence of (3.24) drops out and the only occurrence of M in the above equations is in the expression for E , which contains the term $nN \binom{M}{2} + M|\nu|$. We now define \bar{m} by (3.4) with $m_j^{(a)}$ therein replaced by $m_{N-j}^{(a)}$ (i.e., $\bar{m}_j^{(a)} = m_{N-j}^{(a)}$), and define $\tilde{m}_j^{(a)} = m_{N+j}^{(a)}$. Correspondingly, define \bar{p} ($\tilde{p}_j^{(a)}$) by (3.6) ((3.19)) with m replaced by \bar{m} (\tilde{m}). Then (3.25) ((3.26)) is nothing but (3.6) ((3.19)) with m and p replaced by \bar{m} and \bar{p} (\tilde{m} and \tilde{p}), and (3.24) becomes (after also multiplying by $\bar{\Lambda}_a$ and summing over a)

$$(3.28) \quad \sum_{a=1}^{n-1} \sum_{j=1}^{N-1} j \bar{m}_j^{(a)} \alpha_a = \mu - |\nu| \bar{\Lambda}_1 + N \sum_{a=1}^{n-1} \alpha_a \left[m_N^{(a)} + \sum_{j=1}^{N-1} \bar{m}_j^{(a)} + \sum_{j \geq 1} \tilde{m}_j^{(a)} \right].$$

Now writing the expressions (3.5) and (3.18) without the summations on the left as $F_{\eta\nu; m}(q)$ and $K_{\lambda'\eta; \{m_j^{(a)}\}}(q)$ results in

$$(3.29) \quad q^{-nN \binom{M}{2} - M|\nu|} K_{NM(1^n) + \lambda, (N^{nM}) \cup \nu}(q) \\ = \sum F_{\eta\nu; \bar{m}}(q) K_{\xi'\eta; \{\tilde{m}_j^{(a)}\}}(q) \prod_{a=1}^{n-1} \left[\begin{matrix} m_N^{(a)} + m_a(\eta) + (n-a)M \\ m_a(\eta) \end{matrix} \right]_q,$$

where the sum is over \bar{m} , the $\tilde{m}_j^{(a)}$ and $m_N^{(a)}$ subject to the following restrictions: (i) μ is fixed by (3.27), i.e., by (3.20) with m replaced by \bar{m} but must be an element of P_+ (so that $m_a(\eta) = (\mu|\alpha_a) \geq 0$, and (ii) (3.28) must hold. To understand the occurrence of ξ in the above, note that it is the unique partition such that $|\xi| = |\eta|$ and $\xi_b - |\xi|/n = \lambda_b - |\lambda|/n$.

To complete the proof, replace the above sum by the equivalent sum over $\mu \in P_+$, \bar{m} and the $\tilde{m}_j^{(a)}$ where the latter are subject to the restriction (3.20) with $m \rightarrow \bar{m}$, and \bar{m} is restricted by the condition that (3.28) must fix the $m_N^{(a)}$ to be integers. If we now take the large M limit, the product on the right side of (3.29) yields $1/(q)_{m(\eta)}$ eliminating all $m_N^{(a)}$ -dependence of the summand. Hence we may replace (3.28) by (3.7) with $m \rightarrow \bar{m}$. Finally replacing the sum over μ by a sum over η yields Proposition 3.4.

3.4. Proof of Proposition 3.5. This proposition follows from [15, Props. 4.3, 4.6 and 5.8] and our proof below does not significantly differ from the one given by Hatayama *et al.*

It was shown in [15, Prop. 5.1] that for $|\lambda| = |\mu|$ and $\lambda \in \mathbb{Z}_+^n$ the completely symmetric supernomial can be expressed as

$$\mathcal{S}_{\lambda\mu}(q) = \sum_{\nu} q^{\sum_{a=0}^{n-1} \sum_{j=1}^{\mu_1} (\nu_j^{(a+1)} - \nu_j^{(a)})} \prod_{a=1}^{n-1} \prod_{j=1}^{\mu_1} \left[\begin{matrix} \nu_j^{(a+1)} - \nu_{j+1}^{(a)} \\ \nu_j^{(a+1)} - \nu_j^{(a)} \end{matrix} \right]_q,$$

where the sum is over sequences of partitions $\nu = (\nu^{(1)}, \dots, \nu^{(n-1)})$ such that $\emptyset = \nu^{(0)} \subset \nu^{(1)} \subset \dots \subset \nu^{(n)} = \mu'$ and $|\nu^{(a+1)}| - |\nu^{(a)}| = \lambda_{n-a}$. For $n = 2$ this is equivalent to [43, Eqs. (2.9)–(2.10)].

Now assume that $\mu_1 = N$ and introduce the variables $m_j^{(a)} = \nu_{N-j}^{(n-a)} - \nu_{N-j+1}^{(n-a)}$ for $j \in \{0, \dots, N-1\}$. Also defining the vector m as in (3.4) gives

$$\begin{aligned} \mathcal{S}_{\lambda\mu}(q) &= q^{n(\mu) - \frac{(n-1)|\mu|^2}{2nN} + \frac{1}{2N} \sum_{a=1}^n (\lambda_a^2 - \frac{|\lambda|^2}{n^2})} \\ &\quad \times \sum q^{\frac{1}{2}m(C \otimes \bar{C}^{-1})m - \sum_{i=1}^{l(\mu)} (e_1 \otimes \bar{e}_{N-\mu_i})(I \otimes \bar{C}^{-1})m} \prod_{a=1}^{n-1} \prod_{j=0}^{N-1} \left[\begin{matrix} m_j^{(a)} + p_j^{(a)} \\ m_j^{(a)} \end{matrix} \right]_q, \end{aligned}$$

where the sum is over m and the $m_0^{(a)}$ such that $\sum_{j=0}^{N-1} (N-j)m_j^{(a)} = \sum_{b=a+1}^n \lambda_b$. The auxiliary variables $p_j^{(a)}$ are defined as $p_j^{(a)} = \sum_{k=0}^j (m_k^{(a-1)} - m_k^{(a)}) + \delta_{a,1} \mu'_{N-j}$ (with $m_j^{(0)} := 0$). Making the replacements (3.21) in the above result, one finds

$$\begin{aligned} q^{-nN \binom{M}{2} - M|\nu|} \mathcal{S}_{NM(1^n) + \lambda, (N^n M) \cup \nu}(q) &= q^{n(\nu) - \frac{(n-1)|\nu|^2}{2nN} + \frac{(\mu|\mu)}{2N}} \\ &\quad \times \sum q^{\frac{1}{2}m(C \otimes \bar{C}^{-1})m - \sum_{i=1}^{l(\nu)} (e_1 \otimes \bar{e}_{N-\nu_i})(I \otimes \bar{C}^{-1})m} \prod_{a=1}^{n-1} \prod_{j=0}^{N-1} \left[\begin{matrix} m_j^{(a)} + p_j^{(a)} \\ m_j^{(a)} \end{matrix} \right]_q. \end{aligned}$$

Here λ, ν and μ are as in Proposition 3.5, the sum is over m and the $m_0^{(a)}$ such that

$$(3.30) \quad \sum_{a=1}^{n-1} \sum_{j=1}^{N-1} j m_j^{(a)} \alpha_a = \mu - |\nu| \bar{\Lambda}_1 + N \sum_{a=1}^{n-1} \alpha_a \left[\sum_{j=0}^{N-1} m_j^{(a)} - (n-a)M \right]$$

and $p_j^{(a)} = \sum_{k=0}^j (m_k^{(a-1)} - m_k^{(a)}) + \delta_{a,1} (nM + \nu'_{N-j})$. To arrive at the exponent of q on the right we have used (2.1) and $\sum_{a=1}^n (\lambda_a^2 - |\lambda|^2/n^2) = (\mu|\mu)$, and to obtain the restriction in the sum we have used that $\sum_{a=1}^{n-1} \alpha_a \sum_{b=a+1}^n \lambda_b = |\nu| \bar{\Lambda}_1 - \mu$. Since the exponent of q does not contain M , $m_0^{(a)}$ and $p_j^{(a)}$ it is straightforward to let M tend to infinity. All we need to do is replace $m_0^{(a)} \rightarrow (n-a)M + m_0^{(a)}$. Then the product over the q -binomials becomes

$$P := \prod_{a=1}^{n-1} \left(\left[\begin{matrix} (n-a)M + m_0^{(a)} + p_0^{(a)} \\ (n-a)M + m_0^{(a)} \end{matrix} \right]_q \prod_{j=1}^{N-1} \left[\begin{matrix} m_j^{(a)} + p_j^{(a)} \\ m_j^{(a)} \end{matrix} \right]_q \right),$$

with $p_j^{(a)} = \sum_{k=0}^j (m_k^{(a-1)} - m_k^{(a)}) + M + \delta_{a,1} \nu'_{N-j}$ so that $\lim_{M \rightarrow \infty} P = 1/[(q)_\infty^{n-1}(q)_m]$ which is independent of the $m_0^{(a)}$. Moreover, the term $(n-a)M$ on the right of (3.30) is cancelled so that summing over m and the $m_0^{(a)}$ is equivalent to summing over m subject to the restriction (3.7).

4. BOSONIC REPRESENTATION OF THEOREM 3.1

The aim of this section is to provide an alternative representation (for a special case) of Theorem 3.1, in which γ_k and δ_η are expressed as an $A_{n-1}^{(1)}$ string function and an $A_{n-1}^{(1)}$ configuration sum, both at level N . The advantage of this “bosonic” representation of the theorem is that it suggests a fractional-level generalization in which the integer N is replaced by a rational number.

Let $\Lambda_0, \dots, \Lambda_{n-1}$ be the fundamental weights of the affine Lie algebra $A_{n-1}^{(1)}$ [20], and define Λ_i for all integers i by $\Lambda_i = \Lambda_j$ if $i \equiv j \pmod{n}$. The set of level- N dominant integral weights P_+^N is the set of weights of the form $\Lambda = \sum_{i=0}^{n-1} a_i \Lambda_i$ with $a_i \in \mathbb{Z}_+$ and $\sum_{i=0}^{n-1} a_i = N$. The classical part of a weight $\Lambda \in P_+^N$ will be denoted by $\bar{\Lambda}$, consistent with our definition of the fundamental weights of A_{n-1} . Hence $\bar{\Lambda}_0 = \emptyset$. When $N = 0$, $P_+^N = \emptyset$, but it will be convenient to follow the convention of partitions, and to somewhat ambiguously write $P_+^0 = \{\emptyset\}$, with \emptyset the “empty” level-0 weight. The modular anomaly m_Λ for $\Lambda \in P_+^N$ is given by

$$m_\Lambda = \frac{\|\bar{\Lambda} + \rho\|^2}{2(n+N)} - \frac{\|\rho\|^2}{2n},$$

with $\|\rho\|^2 = n \dim(A_{n-1})/12 = n(n^2 - 1)/12$. We will also employ a shifted modular anomaly

$$\bar{m}_\Lambda = \frac{\|\bar{\Lambda} + \rho\|^2}{2(n+N)}.$$

We now define a level- N , $A_{n-1}^{(1)}$ configuration sum $X_{\eta, \Lambda, \Lambda'}$ by

$$(4.1) \quad X_{\eta, \Lambda, \Lambda'}(q) = q^{-\bar{m}_\Lambda} \sum_{\sigma \in S_n} \epsilon(\sigma) \sum_{k \in (n+N)Q + \sigma(\bar{\Lambda} + \rho)} q^{\frac{(k|k)}{2(n+N)}} \left[\begin{matrix} m(\eta) \\ \frac{|\eta|}{n} (1^n) + k - \Lambda' - \rho \end{matrix} \right]$$

for $\Lambda, \Lambda' \in P_+^N$ and $\eta \in \mathcal{P}$ such that $\eta_1 \leq n-1$ and $|\eta| \bar{\Lambda}_1 - \bar{\Lambda} + \bar{\Lambda}' \in Q$.

Using a sign-reversing involution, Schilling and Shimozono [40] showed that $X_{\eta, \Lambda, N\Lambda_0}(q)$ is a special case of the level- N restricted generalized Kostka polynomial. Equating this with the fermionic representation for this polynomial [41] leads to identities for $X_{\eta, \Lambda, N\Lambda_0}$, which for a special subset of $\Lambda \in P_+^N$ can be stated as follows [40, 41] (see also [38]).

Theorem 4.1. *Let $\Lambda = (N-s)\Lambda_r + s\Lambda_{r+1}$ with $r \in \{0, \dots, n-1\}$ and $s \in \{0, \dots, N\}$ and let $\eta \in \mathcal{P}$ such that $\eta_1 \leq n-1$ and $|\eta| \bar{\Lambda}_1 - \bar{\Lambda} \in Q$. Then*

$$(4.2) \quad X_{\eta, \Lambda, N\Lambda_0}(q) = F_{\eta, (sNr)}(q).$$

Note that the second restriction on η is equivalent to $|\eta| \equiv rN + s \pmod{n}$. Also note that from (3.9), (3.11) and (4.2) follow the particularly simple (to state, *not* to prove) identities [40, Eq. (6.5)]

$$(4.3) \quad X_{\eta, \Lambda_i, \Lambda_0}(q) = q^{\frac{1}{2}(\mu|\mu) - \frac{1}{2}\|\bar{\Lambda}_i\|^2}$$

for $|\eta| \equiv i \pmod{n}$ with $\mu = \sum_{a=1}^{n-1} m_a(\eta) \bar{\Lambda}_a$, and [40, Eq. (6.6)]

$$(4.4) \quad X_{\eta, \emptyset, \emptyset}(q) = \delta_{\eta, \emptyset}$$

for $|\eta| \equiv 0 \pmod{n}$.

Next we shall relate $C_{\mu, (sNr)}$ to the string functions of $A_{n-1}^{(1)}$. For $\lambda \in P$ and m a positive integer, the classical A_{n-1} theta function of degree m and characteristics λ is defined by

$$(4.5) \quad \Theta_{\lambda, m}(x; q) = \sum_{k \in mQ + \lambda} q^{\frac{1}{2m}(k|k)} x^k.$$

According to the Weyl–Kac formula [17, 20] the (normalized) character χ_Λ of the integrable highest weight module $L(\Lambda)$ of highest weight $\Lambda \in P_+^N$ can be expressed in terms of theta functions as

$$\chi_\Lambda(x; q) = \frac{\sum_{\sigma \in S_n} \epsilon(\sigma) \Theta_{\sigma(\bar{\Lambda} + \rho), n+N}(x; q)}{\sum_{\sigma \in S_n} \epsilon(\sigma) \Theta_{\sigma(\rho), n}(x; q)}.$$

The level- N , $A_{n-1}^{(1)}$ string functions $C_{\mu, \Lambda}$ for $\mu \in P$ and $\Lambda \in P_+^N$ arise as coefficients in the expansion of χ_Λ in terms of degree- N theta functions [19]:

$$(4.6) \quad \chi_\Lambda(x; q) = \sum_{\mu \in P/NQ} C_{\mu, \Lambda}(q) \Theta_{\mu, N}(x; q).$$

Two simple properties of the string functions are [21]

$$C_{\mu, \Lambda}(q) = 0 \text{ if } \mu - \bar{\Lambda} \notin Q$$

and

$$(4.7) \quad C_{\sigma(\mu) + N\alpha, \Lambda} = C_{\mu, \Lambda} \text{ for } \alpha \in Q \text{ and } \sigma \in S_n.$$

For our purposes it will be convenient to also introduce normalized string functions $\mathcal{C}_{\mu, \Lambda}$ through

$$\mathcal{C}_{\mu, \Lambda}(q) = q^{\frac{1}{2N}(\mu|\mu) - m_\Lambda} C_{\mu, \Lambda}(q).$$

The relevance of the string functions to $C_{\mu\nu}(q)$ of (3.8) lies in the following theorem.

Theorem 4.2. *Let $\Lambda = (N-s)\Lambda_r + s\Lambda_{r+1}$ with $r \in \{0, \dots, n-1\}$ and $s \in \{0, \dots, N\}$ and let $\mu \in P_+$ such that $\mu - \bar{\Lambda} \in Q$. Then*

$$(4.8) \quad \mathcal{C}_{\mu, \Lambda}(q) = C_{\mu, (sN^r)}(q).$$

This theorem is the culmination of quite a number of results. For $N = 1$ we have according to (3.10)

$$(4.9) \quad \mathcal{C}_{\mu, \Lambda_i}(q) = \frac{q^{\frac{1}{2}(\mu|\mu) - \frac{1}{2}|\bar{\Lambda}_i|^2}}{(q)_\infty^{n-1}}$$

for $\mu - \bar{\Lambda}_i \in Q$. This was first obtained by Kac in [18]. For $n = 2$ but arbitrary N the above theorem is due to Lepowsky and Primc [29] (see also [11]). For general n the theorem was conjectured by Kuniba *et al.* [27] for $r = s = 0$ and proved by Georgiev [14] for $r = 0$ and $r = n - 1$, and by Hatayama *et al.* [15, Prop. 5.8] for general r .

If we now take the conjugate Bailey pair of Theorem 3.1 with $\nu = (sN^r)$ and eliminate $F_{\eta\nu}$ and $C_{\ell\bar{\Lambda}_1 - \pi(k), \nu}$ by virtue of Theorems 4.1 and 4.2, we arrive at the $\Lambda = (N-s)\Lambda_r + s\Lambda_{r+1}$ case of the following conjecture.

Conjecture 4.3. *Let $\ell \in \mathbb{Z}_+$, $k \in Q \cap P_+$ and $\eta \in \mathcal{P}$ such that $\eta_1 \leq n-1$ and $|\eta| \equiv \ell \pmod{n}$. For $\Lambda \in P_+^N$ such that $\ell\bar{\Lambda}_1 - \bar{\Lambda} \in Q$*

$$\gamma_k = \mathcal{C}_{\ell\bar{\Lambda}_1 - \pi(k), \Lambda}(q) \quad \text{and} \quad \delta_\eta = X_{\eta, \Lambda, N\Lambda_0}(q)$$

form an A_{n-1} conjugate Bailey pair relative to q^ℓ .

For $\Lambda = (N-s)\Lambda_r + s\Lambda_{r+1}$ we can of course claim this as a theorem thanks to Theorems 4.1 and 4.2. For $n = 2$ a proof of the above result is implicit in [42]. In the next section it will be shown that Conjecture 4.3 follows by manipulating Corollary 2.2.

5. FRACTIONAL-LEVEL CONJUGATE BAILEY PAIRS

The advantage of Conjecture 4.3 over Theorem 3.1 is that it readily lends itself to further generalization. In order to describe this we extend our definitions of the configuration sum $X_{\eta,\Lambda,\Lambda'}$ and string function $\mathcal{C}_{\mu,\Lambda}$ to fractional levels.

Let p and p' be integers such that $p \geq 1$ and $p' \geq n$, and fix the level N in terms of p and p' as

$$(5.1) \quad N = p'/p - n.$$

In contrast with the previous two sections, $N \in \mathbb{Q}$ with $N > -n$. Given p and p' the set $P^{(p,p')}$ of cardinality $\binom{p'-1}{n-1}$ is defined as the set of weights Λ of the form $\Lambda = \sum_{i=0}^{n-1} a_i \Lambda_i$ such that $\bar{\Lambda} \in P_+$ (i.e., $a_i \in \mathbb{Z}_+$ for $i \geq 1$), $\sum_{i=1}^{n-1} a_i \leq p' - n$ and $\sum_{i=0}^{n-1} a_i = N$. Hence $P_+^{(1,p')} = P_+^N$, but for $p \geq 2$ the coefficient a_0 is noninteger if p is not a divisor of p' ; $p'/p - a_0 \in \{n, n+1, \dots, p'\}$. Of course, whenever we write P_+^N it is assumed that $N \in \mathbb{Z}_+$. Note that the map $\Lambda \rightarrow \Lambda + (1 - 1/p)p'\Lambda_0$ defines a bijection between $P^{(p,p')}$ and $P_+^{p'-n}$. For p and p' relatively prime the set $P^{(p,p')}$ is a special subset of the set of so-called admissible level- N weights [22, 23] in that only those weights are included whose classical part is in P_+ . We do not know how to include more general admissible weights in the results of this section.

The definition of the configuration sum (4.1) is most easily generalized to fractional N , and for $\Lambda, \Lambda' \in P^{(p,p')}$ and $\eta \in \mathcal{P}$ such that $\eta_1 \leq n - 1$ and $|\eta|\bar{\Lambda}_1 - \bar{\Lambda} + \bar{\Lambda}' \in Q$, we define

$$(5.2) \quad X_{\eta,\Lambda,\Lambda'}(q) = q^{-\bar{m}\Lambda} \sum_{\sigma \in S_n} \epsilon(\sigma) \sum_{k \in p'Q + \sigma(\bar{\Lambda} + \rho)} q^{\frac{(k|k)}{2(n+N)}} \left[\frac{|\eta|}{n} (1^n) + k - \bar{\Lambda}' - \rho \right].$$

If $|\eta|\bar{\Lambda}_1 - \bar{\Lambda} + \bar{\Lambda}' \notin Q$ we set $X_{\eta,\Lambda,\Lambda'}(q) = 0$.

An important remark concerning (5.2) is in order. We have not imposed that p and p' be relatively prime, and as a consequence, N does not uniquely fix p and p' . Hence, given an admissible weight of level N , one cannot determine p and p' . Put differently, the sets $P^{(p,p')}$ and $P^{(t,t')}$ with $(p,p') \neq (t,t')$ are not necessarily disjoint. For example, $P^{(p,p')} \subseteq P^{(kp, kp')}$ for k a positive integer. As a result of this, definition (5.2) is ambiguous since the right-hand side depends not only on N (which is fixed uniquely by Λ) but also on p' . We however trust that by writing $\Lambda \in P^{(p,p')}$ the reader will have no trouble interpreting (5.2). Those unwilling to accept the above notation may from now on assume that p and p' are relatively prime, or should add a superscript p' or even (p,p') to $X_{\eta,\Lambda,\Lambda'}$.

Before we show how to also generalize (4.6) to yield fractional-level string functions, we shall list several important properties of $X_{\eta,\Lambda,\Lambda'}$, a number of which are conjectural.

Lemma 5.1. *For $\Lambda, \Lambda' \in P^{(p,p')}$, $X_{\emptyset,\Lambda,\Lambda'}(q) = \delta_{\Lambda,\Lambda'}$.*

Proof. From (2.22) or (2.25) it follows that $[\emptyset] = \delta_{\lambda,\emptyset}$. Hence

$$X_{\emptyset,\Lambda,\Lambda'}(q) = q^{\bar{m}\Lambda' - \bar{m}\Lambda} \sum_{\sigma \in S_n} \epsilon(\sigma) \sum_{k \in p'Q + \sigma(\bar{\Lambda} + \rho)} \delta_{k,\bar{\Lambda}' + \rho},$$

where we have used that $q^{(k|k)/(2(n+N))} \delta_{k,\bar{\Lambda}' + \rho} = q^{\bar{m}\Lambda'} \delta_{k,\bar{\Lambda}' + \rho}$. The question now is: can we satisfy $\bar{\Lambda}' + \rho \in p'Q + \sigma(\bar{\Lambda} + \rho)$? Since $\Lambda, \Lambda' \in P^{(p,p')}$ this simplifies to the problem of finding for which Λ, Λ' and σ there holds $\bar{\Lambda}' + \rho = \sigma(\bar{\Lambda} + \rho)$. Since $\bar{\Lambda}' + \rho \in P_+$ and $\sigma(\bar{\Lambda} + \rho) \in P_+$ iff $\sigma = (1, 2, \dots, n)$, the only solution is given by $\Lambda = \Lambda'$ and $\sigma = (1, 2, \dots, n)$, establishing the claim of the lemma. \square

For $\Lambda = \sum_{i=0}^{n-1} a_i \Lambda_i \in P^{(p,p')}$ define $\Lambda^c \in P^{(p,p')}$ as $\Lambda^c = \sum_{i=0}^{n-1} a_i \Lambda_{n-i}$, and for $\eta = (1^{\zeta_1}, 2^{\zeta_2}, \dots, (n-1)^{\zeta_{n-1}})$ define $\eta^c = (1^{\zeta_{n-1}}, 2^{\zeta_{n-2}}, \dots, (n-1)^{\zeta_1})$. Note that the partition

η^c is the complement of η with respect to $(n^{l(\eta)})$, and in our earlier notation of Section 2.1, $\eta^c = \tilde{\eta}_{(n^{l(\eta)})}$.

Lemma 5.2. For $\Lambda, \Lambda' \in P^{(p,p')}$ and $\eta \in \mathcal{P}$ such that $\eta_1 \leq n-1$,

$$(5.3) \quad X_{\eta, \Lambda, \Lambda'} = X_{\eta^c, \Lambda^c, (\Lambda')^c}.$$

Observe that the condition $|\eta| \bar{\Lambda}_1 - \bar{\Lambda} + \bar{\Lambda}' \in Q$ implies that $|\eta^c| \bar{\Lambda}_1 - \bar{\Lambda}^c + (\bar{\Lambda}')^c \in Q$ as it should.

Proof. By the symmetry (2.25) of the antisymmetric supernomials

$$X_{\eta, \Lambda, \Lambda'}(q) = q^{-\bar{m}_\Lambda} \sum_{\sigma \in S_n} \epsilon(\sigma) \sum_{k \in p'Q + \sigma(\bar{\Lambda} + \rho)} q^{\frac{(k|k)}{2(n+N)}} \left[\frac{|\eta^c|}{n} (1^n) - k + \Lambda' + \rho \right].$$

Next use (2.24) with $\sigma = \pi$ and replace k by $-\pi(k)$. Then

$$X_{\eta, \Lambda, \Lambda'}(q) = q^{-\bar{m}_\Lambda} \sum_{\sigma \in S_n} \epsilon(\sigma) \sum_{k \in p'Q - \pi \circ \sigma(\bar{\Lambda} + \rho)} q^{\frac{(k|k)}{2(n+N)}} \left[\frac{|\eta^c|}{n} (1^n) + k + \pi(\Lambda' + \rho) \right].$$

Finally use $\bar{\Lambda} + \rho = -\pi(\bar{\Lambda}^c + \rho)$ (which implies $\bar{m}_\Lambda = \bar{m}_{\Lambda^c}$), $\pi(\bar{\Lambda}' + \rho) = -(\bar{\Lambda}')^c - \rho$ and replace σ by $\pi \circ \sigma \circ \pi$ to find (5.3). \square

Based on extensive computer assisted experiments we are led to the following conjectures.

Conjecture 5.3. Let $p' - p - n + 1 \geq 0$. For $\Lambda, \Lambda' \in P^{(p,p')}$ such that $\Lambda' + (p-1)(p'/p-1)\Lambda_0 \in P_+^{p'-p-n+1}$, $X_{\eta, \Lambda, \Lambda'}(q)$ is a polynomial with nonnegative coefficients.

Conjecture 5.4. For $\Lambda, \Lambda' \in P^{(p,p')}$, $\lim_{|\eta| \rightarrow \infty} X_{\eta, \Lambda, \Lambda'}(q) = 0$.

For $n = 2$ both conjectures readily follow from results in [46]. The last conjecture will be crucial later, ensuring the convergence of certain sums over $X_{\eta, \Lambda, \Lambda'}(q)$.

Next we come to the definition of the fractional-level string functions. For $\Lambda \in P^{(p,p')}$ define

$$(5.4) \quad \chi_\Lambda(x; q) = \frac{\sum_{\sigma \in S_n} \epsilon(\sigma) \Theta_{\sigma(\bar{\Lambda} + \rho), p'}(x; q^p)}{\sum_{\sigma \in S_n} \epsilon(\sigma) \Theta_{\sigma(\rho), n}(x; q)}.$$

For p and p' relatively prime $\chi_\Lambda(x; q)$ is a character of a (special) admissible representation of $A_{n-1}^{(1)}$, (5.4) corresponding to (a special case of) the Kac–Wakimoto character formula for admissible characters [22]. The corresponding string functions at fractional-level N are defined via the formal expansion

$$(5.5) \quad \chi_\Lambda(x; q) = \sum_{\mu \in P} q^{\frac{1}{2N}(\mu|\mu)} C_{\mu, \Lambda}(q) x^\mu = q^{m_\Lambda} \sum_{\mu \in P} C_{\mu, \Lambda}(q) x^\mu$$

for $\Lambda \in P^{(p,p')}$ with N given by (5.1). To see this is consistent with (4.6), note that for $\Lambda \in P_+^N$

$$\begin{aligned} \sum_{\mu \in P} q^{\frac{1}{2N}(\mu|\mu)} C_{\mu, \Lambda}(q) x^\mu &= \sum_{\mu \in P} \sum_{\substack{\nu \in P/NQ \\ \nu \equiv \mu \pmod{NQ}}} q^{\frac{1}{2N}(\mu|\mu)} C_{\nu, \Lambda}(q) x^\mu \\ &= \sum_{\nu \in P/NQ} C_{\nu, \Lambda}(q) \sum_{\mu \in NQ + \nu} q^{\frac{1}{2N}(\mu|\mu)} x^\mu = \sum_{\nu \in P/NQ} C_{\nu, \Lambda}(q) \Theta_{\nu, N}(x), \end{aligned}$$

where the first equality follows from (4.7) with $\sigma = (1, 2, \dots, n)$. Since the string functions for non-integral levels only satisfy (4.7) with $\alpha = \emptyset$, it is generally not possible to rewrite (5.5) as (4.6).

Having defined the configuration sums and string functions for fractional levels we can now state the following generalization of Conjecture 4.3.

Theorem 5.5. *If Corollary 2.2 is true then Conjecture 4.3 holds for all $\Lambda \in P^{(p,p')}$.*

This theorem claims a conjugate Bailey pair relative to q^ℓ with $\delta_\eta = X_{\eta,\Lambda,N\Lambda_0}(q)$. It is not clear a priori that substituting this δ_η in (3.2) will lead to a converging sum, and it is here that Conjecture 5.4 is crucial.

In our proof below we will show that Conjecture 4.3 may be manipulated to yield the following expression for admissible characters of $\Lambda_{n-1}^{(1)}$:

$$(5.6) \quad \chi_\Lambda(q) = q^{m_\Lambda} \sum_{\substack{\mu \in P \\ \mu - \bar{\Lambda} \in Q}} x^\mu \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \bar{\Lambda}_1 - \mu \in Q}} \frac{X_{\eta,\Lambda,N\Lambda_0}(q)}{(q)_{m(\eta)}} \left[\frac{m(\eta)}{n} (1^n) + \mu \right]$$

for $\Lambda \in P^{(p,p')}$. Comparing this with (5.5) leads to

$$(5.7) \quad \mathcal{C}_{\mu,\Lambda}(q) = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \bar{\Lambda}_1 - \mu \in Q}} \frac{X_{\eta,\Lambda,N\Lambda_0}(q)}{(q)_{m(\eta)}} \left[\frac{m(\eta)}{n} (1^n) + \mu \right],$$

for $\mu - \bar{\Lambda} \in Q$ and zero otherwise, from which Theorem 5.5 easily follows.

Proof of Theorem 5.5. Take Corollary 2.2 and expand the Kostka polynomial therein using (2.26). By $\tau(\delta) - \delta = \tau(\rho) - \rho$ this yields

$$\begin{aligned} & \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \equiv |\mu| \pmod{n}}} \sum_{\sigma, \tau \in S_n} \sum_{\lambda \in nQ + \sigma(\rho) - \rho} \epsilon(\sigma) \epsilon(\tau) q^{\frac{1}{2n}(\lambda|\lambda + 2\rho)} \\ & \quad \times \frac{1}{(q)_{m(\eta)}} \left[\frac{m(\eta)}{n} (1^n) + \tau(\nu + \rho) - \rho \right] \left[\frac{m(\eta)}{n} (1^n) + \mu - \lambda \right] = \delta_{\mu,\nu}, \end{aligned}$$

where $\mu, \nu \in \mathcal{P}$ such that $l(\mu), l(\nu) \leq n$ and $|\mu| = |\nu|$.

Next replace μ, ν by $\hat{\mu}, \hat{\nu} \in P_+$ via $\hat{\omega} = \omega - |\omega|(1^n)/n$, so that $\hat{\mu} - \hat{\nu} \in Q$. Making these replacements in the actual equation is trivial and it only needs to be remarked that the restriction $|\eta| \equiv |\mu| \pmod{n}$ becomes $|\eta| \bar{\Lambda}_1 - \hat{\mu} \in Q$. After dropping the hats from $\hat{\mu}$ and $\hat{\nu}$ we thus find

$$(5.8) \quad \begin{aligned} & \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \bar{\Lambda}_1 - \mu \in Q}} \sum_{\sigma, \tau \in S_n} \sum_{\lambda \in nQ + \sigma(\rho) - \rho} \epsilon(\sigma) \epsilon(\tau) q^{\frac{1}{2n}(\lambda|\lambda + 2\rho)} \\ & \quad \times \frac{1}{(q)_{m(\eta)}} \left[\frac{m(\eta)}{n} (1^n) + \tau(\nu + \rho) - \rho \right] \left[\frac{m(\eta)}{n} (1^n) + \mu - \lambda \right] = \delta_{\mu,\nu}, \end{aligned}$$

where $\mu, \nu \in P_+$ such that $\mu - \nu \in Q$. Next we will show that if these conditions are relaxed to $\mu, \nu \in P$ such that $\mu - \nu \in Q$, then the right-hand side of (5.8) needs to be replaced by

$$(5.9) \quad \sum_{\sigma \in S_n} \epsilon(\sigma) \delta_{\mu, \sigma(\nu + \rho) - \rho}.$$

In the following we refer to the identity obtained by equating the left-side of (5.8) with (5.9) (for $\nu, \mu \in P$ such that $\mu - \nu \in Q$) by (I). Now there are three possibilities for $\mu, \nu \in P$: (i) there exist $w, v \in S_n$ such that $w(\mu + \rho) - \rho, v(\nu + \rho) - \rho \in P_+$, (ii) there exists no such w , (iii) there exists no such v . Of course, when w and v exist they are unique and w (v) does not exist if and only if not all components of $w + \rho$ ($v + \rho$) are distinct.

First assume case (i) and set $\mu = w^{-1}(\hat{\mu} + \rho) - \rho$, $\nu = v^{-1}(\hat{\nu} + \rho) - \rho$ with $\hat{\mu}, \hat{\nu} \in P_+$ and replace μ and ν in favour of $\hat{\mu}$ and $\hat{\nu}$. Then change $\lambda \rightarrow w^{-1}(\lambda + \rho) - \rho$, $\sigma \rightarrow w^{-1} \circ \sigma$ and

$\tau \rightarrow v \circ \tau$ on the left and use the symmetry (2.24). On the right change $\sigma \rightarrow w^{-1} \circ \sigma \circ v$. Finally drop the hats of $\hat{\mu}$ and $\hat{\nu}$. The result of all this is that (I) is transformed into itself but now $\mu, \nu \in P_+$. For such μ and ν the only contributing term in the sum (5.9) corresponds to $\sigma = (1, 2, \dots, n)$ and we are back at (5.8). Consequently (I) is true in case (i). Next assume (ii) and replace $\lambda \rightarrow \sigma(\lambda + \rho) - \rho$ on the left and $\sigma \rightarrow \sigma^{-1}$ on both sides of (I). After also using (2.5), the left side then contains the sum

$$\sum_{\sigma \in S_n} \epsilon(\sigma) \left[\begin{matrix} m(\eta) \\ \frac{|\eta|}{n}(1^n) + \sigma(\mu + \rho) - \rho - \lambda \end{matrix} \right],$$

whereas the right side reads $\sum_{\sigma \in S_n} \epsilon(\sigma) \delta_{\sigma(\mu + \rho) - \rho, \nu}$. Since not all components of $\mu + \rho$ are distinct this implies that both sides vanish. Hence (II) is also true in case (ii). Similarly it follows that (I) is true in case (iii), with both sides again vanishing.

To proceed, take identity (I), multiply both sides by $x^{\mu + \rho}$, sum μ over P subject to the restriction $\mu - \nu \in Q$ and finally shift $\nu \rightarrow \nu - \rho$. Adopting definition (2.2) for $\nu \in P$ this leads to

$$\begin{aligned} \sum_{\substack{\mu \in P \\ \mu - \nu + \rho \in Q}} x^{\mu + \rho} \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \bar{\Lambda}_1 - \mu \in Q}} \sum_{\sigma, \tau \in S_n} \sum_{\lambda \in nQ + \sigma(\rho) - \rho} \epsilon(\sigma) \epsilon(\tau) q^{\frac{1}{2n}(\lambda|\lambda + 2\rho)} \\ \times \frac{1}{(q)_{m(\eta)}} \left[\begin{matrix} m(\eta) \\ \frac{|\eta|}{n}(1^n) + \tau(\nu) - \rho \end{matrix} \right] \left[\begin{matrix} m(\eta) \\ \frac{|\eta|}{n}(1^n) + \mu - \lambda \end{matrix} \right] = a_\nu(x), \end{aligned}$$

where $\nu \in P$. We now interchange the sums over μ and λ and replace $\mu \rightarrow \mu + \lambda$ followed by $\lambda \rightarrow \lambda - \rho$. The left side then factorizes into a double sum over σ and λ and a triple sum over μ, η and τ . Recalling the theta function (4.5), the double sum can be recognized as the denominator of the right side of (5.4), which will be abbreviated here to $A_\rho(x; q)$. Hence

$$\begin{aligned} q^{-\frac{1}{2n}\|\rho\|^2} A_\rho(x; q) \sum_{\substack{\mu \in P \\ \mu - \nu + \rho \in Q}} x^\mu \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \bar{\Lambda}_1 - \mu \in Q}} \sum_{\tau \in S_n} \epsilon(\tau) \\ \times \frac{1}{(q)_{m(\eta)}} \left[\begin{matrix} m(\eta) \\ \frac{|\eta|}{n}(1^n) + \tau(\nu) - \rho \end{matrix} \right] \left[\begin{matrix} m(\eta) \\ \frac{|\eta|}{n}(1^n) + \mu \end{matrix} \right] = a_\nu(x). \end{aligned}$$

Now fix integers $p \geq 1$ and $p' \geq n$, define N by (5.1) and let $\Lambda \in P^{(p, p')}$. Then multiply both sides of the above equation by $q^{(\nu|\nu)/(2(n+N))}$ and sum over ν such that $\nu \in p'Q + \bar{\Lambda} + \rho$. On the left swap the sum over ν with the other three sums and replace $\nu \rightarrow \tau^{-1}(\nu)$. Recalling the Kac–Wakimoto character formula (5.4) and the definition (5.2) of the configuration sum $X_{\eta, \Lambda, \Lambda'}$, this yields (5.6) and thus (5.7). To complete the proof choose $\mu = \ell \bar{\Lambda}_1 - \pi(k)$, and use (2.24) with $\sigma = \pi$ and $\pi(\ell \bar{\Lambda}_1 - \pi(k)) = -k - \ell \Lambda_{n-1}$. Then (5.7) transforms into the claim of the theorem. \square

6. APPLICATIONS

In this final section we give several applications of our A_{n-1} Bailey lemma, resulting in some new A-type q -series identities. Following the derivation of the Rogers–Ramanujan identities outlined in the introduction, we should try to find pairs of sequences (α, β) that satisfy the defining relation (3.1) of a Bailey pair. If these, together with the A_{n-1} conjugate Bailey pairs of Theorem 3.1 or 5.5, are substituted in (3.3) some hopefully interesting q -series identities will result.

According to (3.1), what is needed to obtain Bailey pairs are polynomial identities involving the antisymmetric supernomials. Examples of such identities, from which Bailey pairs may

indeed be extracted, are (4.2)–(4.4). To however explicitly write down the Bailey pairs arising from these identities is rather cumbersome due to the fact that we first have to apply

$$\sum_{k \in P} f_k = \sum_{k \in P_+} \sum_{\tau \in S_n / S_n^k} f_{\tau(k)}$$

to rewrite the sum over k in (4.1) as a sum over $k \in Q \cap P_+$. Moreover, once the substitutions in (3.3) have been made, one invariably wants to simplify the resulting identity by writing the sum over $k \in Q \cap P_+$ on the left as a sum over $k \in Q$. In order to eliminate the undesirable intermediate step where all sums are restricted to $k \in Q \cap P_+$, we will never explicitly write down Bailey pairs, but only work with the polynomial identities that imply these pairs. To illustrate this in the case of the first Rogers–Ramanujan identity, let us take the Bailey pair (1.6) for $a = 1$ and substitute this in the defining relation (1.1) of an A_1 Bailey pair. After a renaming of variables this leads to the polynomial identity

$$(6.1) \quad \sum_{j=-L}^L (-1)^j q^{j(3j-1)/2} \left[\begin{matrix} 2L \\ L-j \end{matrix} \right]_q = \frac{(q)_{2L}}{(q)_L}.$$

Next take the conjugate Bailey pair (1.4) for $a = 1$ and substitute this into (1.2) to find

$$(6.2) \quad \sum_{r=j}^{\infty} \frac{q^{r^2}}{(q)_{2r}} \left[\begin{matrix} 2r \\ r-j \end{matrix} \right]_q = \frac{q^{j^2}}{(q)_{\infty}}$$

for $j \in \mathbb{Z}_+$. Since both sides are even functions of j we may assume $j \in \mathbb{Z}$. Now multiply (6.2) by $(-1)^j q^{j(3j-1)/2}$ and sum j over the integers. After a change in the order of summation, (6.1) may be applied to carry out the j -sum on the left, yielding

$$\sum_{r=0}^{\infty} \frac{q^{r^2}}{(q)_r} = \frac{1}{(q)_{\infty}} \sum_{j=-\infty}^{\infty} (-1)^j q^{j(5j-1)/2}.$$

This is exactly the identity obtained when (1.6) for $a = 1$ is substituted into (1.5).

In the following we will mimic the above manipulations, albeit in a more general setting. First, if we substitute the conjugate Bailey pair of Theorem 5.5 in the defining relation (3.2) and generalize the symmetrization immediately after (6.2), we arrive back at (5.7) which shall therefore serve as our starting point. Now let p, p', t, t' be integers such that $p, t \geq 1$ and $p', t' \geq n$, and set $N = p'/p - n$ and $M = t'/t - n$. Take (5.7) with $\Lambda \in P^{(t, t')}$, replace μ by $k - \rho - \Lambda''$ and multiply both sides by $\epsilon(\sigma) q^{-\bar{m}_{\Lambda'} + (k|k)/(2(n+N))}$ where $\Lambda', \Lambda'' \in P^{(p, p')}$ and $\sigma \in S_n$. Then sum over k such that $k \in p'Q + \sigma(\bar{\Lambda}' + \rho)$ and sum σ over the elements of S_n . Using the definition (5.2) of $X_{\eta, \Lambda', \Lambda''}$ this yields

$$(6.3) \quad b_{\Lambda, \Lambda', \Lambda''}(q) = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta|_{\bar{\Lambda}' - \bar{\Lambda}} \in Q}} \frac{X_{\eta, \Lambda, M\Lambda_0}(q) X_{\eta, \Lambda', \Lambda''}(q)}{(q)_{m(\eta)}}$$

with $b_{\Lambda, \Lambda', \Lambda''}$ defined as

$$(6.4) \quad b_{\Lambda, \Lambda', \Lambda''}(q) = q^{-\bar{m}_{\Lambda'}} \sum_{\sigma \in S_n} \epsilon(\sigma) \sum_{k \in p'Q + \sigma(\bar{\Lambda}' + \rho)} q^{\frac{(k|k)}{2(n+N)}} C_{k - \bar{\Lambda}'' - \rho, \Lambda}(q)$$

and $\Lambda \in P^{(t, t')}$, $\Lambda', \Lambda'' \in P^{(p, p')}$ such that $\bar{\Lambda} - \bar{\Lambda}' + \bar{\Lambda}'' \in Q$.

It is interesting to note that (6.3) implies the highly nontrivial symmetry

$$(6.5) \quad b_{\Lambda, \Lambda', N\Lambda_0} = b_{\Lambda', \Lambda, M\Lambda_0}.$$

Another symmetry, which follows from (6.3), (5.3) and $(q)_{m(\eta)} = (q)_{m(\eta^c)}$ is

$$(6.6) \quad b_{\Lambda, \Lambda', \Lambda''} = b_{\Lambda^c, (\Lambda')^c, (\Lambda'')^c}.$$

Unlike (6.5), however, this also follows from (6.4). Since $\bar{\Lambda} + \rho = -\pi(\bar{\Lambda}^c + \rho)$ it follows from (5.4) that $\chi_{\Lambda}(x; q) = \chi_{\Lambda^c}(1/x; q)$ and hence that $\mathcal{C}_{\mu, \Lambda} = \mathcal{C}_{-\mu, \Lambda^c}$. Using this in (6.4), replacing $k \rightarrow -\pi(k)$ and $\sigma \rightarrow \pi \circ \sigma \circ \pi$ and using $\mathcal{C}_{\mu, \Lambda} = \mathcal{C}_{\pi(\mu), \Lambda}$, (6.6) once more follows.

When $\Lambda \in P_+^M$, (6.4) coincides with a special case of an expression of Kac and Wakimoto for the coset branching function $b_{\Lambda''}^{\Lambda \otimes \Lambda'}$ defined by

$$\chi_{\Lambda}(x; q)\chi_{\Lambda'}(x; q) = \sum_{\substack{\Lambda'' \in P^{(p, p' + Mp)} \\ \bar{\Lambda} + \bar{\Lambda}' - \bar{\Lambda}'' \in Q}} q^{\phi_{\Lambda, \Lambda', \Lambda''}} b_{\Lambda''}^{\Lambda \otimes \Lambda'}(q)\chi_{\Lambda''}(x; q)$$

for $\Lambda \in P_+^M$, $\Lambda' \in P^{(p, p')}$ and

$$\phi_{\Lambda, \Lambda', \Lambda''} = \frac{\|(p' + Mp)(\bar{\Lambda}' + \rho) - p'(\bar{\Lambda}'' + \rho)\|^2}{2Mp'(p' + Mp)} - \frac{\|\bar{\Lambda}' - \bar{\Lambda}''\|^2}{2M} + m_{\Lambda}.$$

According to [23, Eq. (3.1.1)],

$$b_{\Lambda''}^{\Lambda \otimes \Lambda'}(q) = q^{-\bar{m}_{\Lambda'}} \sum_{\sigma \in S_n} \epsilon(\sigma) \sum_{k \in p'Q + \sigma(\bar{\Lambda}' + \rho)} q^{\frac{(k|k)}{2(n+N)}} \mathcal{C}_{\bar{\Lambda}'' + \rho - k, \Lambda}(q).$$

Recalling that $\mathcal{C}_{\mu, \Lambda} = \mathcal{C}_{-\mu, \Lambda^c}$ and comparing the expression for the branching function with (6.4) leads to the identification

$$(6.7) \quad b_{\Lambda, \Lambda', \Lambda''} = b_{M\Lambda_0 + \Lambda''}^{\Lambda^c \otimes \Lambda'}$$

for $\Lambda \in P_+^M$ and $\Lambda', \Lambda'' \in P^{(p, p')}$. Note in particular that $M\Lambda_0 + \Lambda'' \in P^{(p, p' + Mp)}$ and $\bar{\Lambda} - \bar{\Lambda}' + \bar{\Lambda}'' \in Q \Leftrightarrow \bar{\Lambda}^c + \bar{\Lambda}' - \bar{\Lambda}'' \in Q$ as it should.

Thus far we have only manipulated the conjugate Bailey pair of Theorem 5.5, and to turn (6.3) into more explicit q -series identities we need to specify (p, p') and/or (t, t') such that the configuration sums on the right can be evaluated in closed form. As a first example choose $(p, p') = (1, n)$. Then $\Lambda', \Lambda'' \in P_+^0$ and hence $\bar{\Lambda}' = \bar{\Lambda}'' = \emptyset$. Recalling (4.4) and Lemma 5.1 it follows that the right side of (6.3) trivializes to $\delta_{\Lambda, M\Lambda_0}$. Making the change $k \rightarrow \sigma(nk + \rho)$ in (6.4) and using $\mathcal{C}_{\mu, \Lambda} = \mathcal{C}_{\sigma(\mu), \Lambda}$ then yields

$$(6.8) \quad \sum_{k \in Q} q^{\frac{1}{2}(k|nk+2\rho)} \sum_{\sigma \in S_n} \epsilon(\sigma) \mathcal{C}_{nk+\rho-\sigma(\rho), \Lambda}(q) = \delta_{\Lambda, M\Lambda_0}$$

for $\Lambda \in P^{(t, t')}$ such that $\bar{\Lambda} \in Q$. This is an A_{n-1} , fractional-level analogue of Euler's pentagonal number theorem. If for $\Lambda \in P_+^1$ we use (4.9) and perform the sum over σ by the Vandermonde determinant (2.3) in the form

$$(6.9) \quad \sum_{\sigma \in S_n} \epsilon(\sigma) x^{\sigma(\rho) - \rho} = \prod_{1 \leq i < j \leq n} (1 - x_j/x_i).$$

with $x_i = q^{-nk_i - n + i}$, we find

$$\sum_{k \in Q} q^{\frac{1}{2}n(n+1)(k|k) + (k|\rho)} \prod_{1 \leq i < j \leq n} (1 - q^{n(k_i - k_j) + j - i}) = (q)_{\infty}^{n-1}.$$

For $n = 2$ this is Euler's famous sum

$$\sum_{k=-\infty}^{\infty} (-1)^k q^{k(3k+1)/2} = (q)_{\infty},$$

and for general n it corresponds to a specialization of the A_{n-1} Macdonald identity [31]. For $\Lambda \in P_+^N$ identity (6.8) is [23, Eq. (2.1.17)]; $\lambda = k\Lambda_0$ of Kac and Wakimoto, and for $\Lambda \in P^{(t, t')}$ with $n = 2$ it is [46, Prop 8.2; $\eta = 0$].

Many more identities arise if we evoke Theorem 4.1. In fact, instead of just this theorem we will apply the following generalization which assumes the notation of (3.5).

Conjecture 6.1. *Let $\eta \in \mathcal{P}$ such that $\eta_1 \leq n-1$. For $s, u \in \{0, \dots, N\}$, $r, v \in \{0, \dots, n-1\}$ let $\Lambda = (N-s)\Lambda_r + s\Lambda_{r+1}$ and $\Lambda' = (N-u)\Lambda_0 + u\Lambda_{n-v}$ such that $|\eta|\bar{\Lambda}_1 - \bar{\Lambda} + \bar{\Lambda}' \in Q$ ($|\eta| \equiv rN + s + uv \pmod{n}$), and define*

$$F_{\eta, \Lambda, \Lambda'}(q) = q^{\frac{\|\mu - \bar{\Lambda}'\|^2 - \|\bar{\Lambda}\|^2}{2N}} \sum_m q^{\frac{1}{2}m(C \otimes \bar{C}^{-1})m - (e_1 \otimes \bar{e}_{N-s})(I \otimes \bar{C}^{-1})m} \begin{bmatrix} m+p \\ m \end{bmatrix}_q,$$

where p is determined by

$$(C \otimes \bar{I})m + (I \otimes \bar{C})p = (m(\eta) \otimes \bar{e}_1) + (e_1 \otimes \bar{e}_{N-s}) + (e_v \otimes \bar{e}_{N-u})$$

and the sum is over m such that

$$\sum_{a=1}^{n-1} \sum_{j=1}^{N-1} jm_j^{(a)} \alpha_a \in NQ + \mu - (rN + s)\bar{\Lambda}_1 - u\bar{\Lambda}_v$$

with $\mu = \sum_{a=1}^{n-1} m_a(\eta)\bar{\Lambda}_a \in P_+$. Then

$$X_{\eta, \Lambda, \Lambda'}(q) = F_{\eta, \Lambda, \Lambda'}(q).$$

With Λ as given above, $F_{\eta, \Lambda, N\Lambda_0} = F_{\eta, (sN)r}$ and in this case the conjecture becomes Theorem 4.1. For $n=2$ Conjecture 6.1 follows from polynomial identities proved in [8, 51].

From (6.3), (6.7) and Conjecture 6.1 we infer the following fermionic representation for the branching functions:

$$b_{M\Lambda_0 + \Lambda''}^{\Lambda^c \otimes \Lambda'}(q) = b_{M\Lambda_0 + (\Lambda'')^c}^{\Lambda \otimes (\Lambda')^c}(q) = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \equiv bM+a \pmod{n}}} \frac{F_{\eta, \Lambda, M\Lambda_0}(q) F_{\eta, \Lambda', \Lambda''}(q)}{(q)_{m(\eta)}}$$

with $\Lambda = (M-a)\Lambda_b + a\Lambda_{b+1}$, $\Lambda' = (N-s)\Lambda_r + s\Lambda_{r+1}$ and $\Lambda'' = (N-u)\Lambda_0 + u\Lambda_{n-v}$ such that $\bar{\Lambda} - \bar{\Lambda}' + \bar{\Lambda}'' \in Q$ (i.e., $bM + a \equiv rN + s + uv \pmod{n}$). Here $b, r, v \in \{0, \dots, n-1\}$, $a \in \{0, \dots, M\}$ and $s, u \in \{0, \dots, N\}$. Note that for $\Lambda'' = N\Lambda_0$ this result is certainly true thanks to Theorem 4.1. For $b = r = s = 0$ (so that $a \equiv uv \pmod{n}$) the above identity has also been obtained by Schilling and Shimozono [41, Eq. (7.10)] who exploited the fact that

$$b_{(M+N-u)\Lambda_0 + u\Lambda_v}^{(M-a)\Lambda_0 + a\Lambda_1 \otimes N\Lambda_0}(q) = \lim_{L \rightarrow \infty} q^{-akL - nM \binom{kL}{2}} K_{\lambda^{(L)}, R^{(L)}}^{N+M}(q).$$

Here $K_{\lambda, R}^N(q)$ is the level-restricted generalized Kostka polynomial, $k \in [n-1]$, $\lambda^{(L)} = (c^{n-v}(c+u)^v)$ with $c = kLM + (a-uv)/n$, and $R^{(L)} = (R_1, \dots, R_{L_{n+1}})$, with $R_1 = (a)$ and $R_i = (M^k)$ for $i \geq 2$. By using the generalization of Theorem 4.1 to all $\Lambda \in P_+^N$ [38, 40, 41] we can also apply (6.3) to prove fermionic formulae for all of the branching functions $b_{(N+M)\Lambda_0}^{\Lambda \otimes \Lambda'}$ with $\Lambda \in P_+^M$ and $\Lambda' \in P_+^N$.

Another nice specialization of (6.3) arises for $\Lambda \in P_+^1$ and $\Lambda'' = N\Lambda_0$. Then $\bar{\Lambda} - \bar{\Lambda}' \in Q$ fixes Λ , and by (4.3), (4.9) and (6.9) it follows that all dependence on Λ cancels out. Dropping the prime in Λ' , this yields

$$(6.10) \quad \frac{q^{\frac{1}{2}\|\bar{\Lambda}\|^2}}{(q)_{\infty}^{n-1}} \sum_{k \in Q} q^{\frac{1}{2}p'(p+p')(k|k) + (p+p')(k|\bar{\Lambda}) + p(k|\rho)} \\ \times \prod_{1 \leq i < j \leq n} (1 - q^{p'(k_i - k_j) + j - i + (\bar{\Lambda}|\varepsilon_i - \varepsilon_j)}) = \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta|\bar{\Lambda}_1 - \bar{\Lambda} \in Q}} \frac{q^{\frac{1}{2}(\mu|\mu)} X_{\eta, \Lambda, N\Lambda_0}(q)}{(q)_{m(\eta)}}$$

for $\Lambda \in P^{(p,p')}$ and $\mu = \sum_{a=1}^{n-1} m_a(\eta) \bar{\Lambda}_a$. When $p' = n$ the sum on the left can be performed by the A_{n-1} Macdonald identity, resulting in

$$(6.11) \quad \sum_{\substack{\eta \in \mathcal{P} \\ \eta_1 \leq n-1 \\ |\eta| \equiv 0 \pmod{n}}} \frac{q^{\frac{1}{2}(\mu|\mu)} X_\eta^{p,n}(q)}{(q)_{m(\eta)}} = \frac{(q^{n+p}; q^{n+p})_\infty^{n-1}}{(q)_\infty^{n-1}} \prod_{i=1}^{n-1} (q^i, q^{n+p-i}; q^{n+p})_\infty^{n-i}$$

with

$$X_\eta^{p,n}(q) = \sum_{\sigma \in S_n} \epsilon(\sigma) \sum_{k \in nQ} q^{\frac{p}{2n}(k|k+2\rho)} \left[\frac{m(\eta)}{n} (1^n) + k + \rho - \sigma(\rho) \right].$$

Here $X_\eta^{p,n}(q) = X_{\eta, N\Lambda_0, N\Lambda_0}(q)$ for $N = n(1/p - 1)$ and $N\Lambda_0 \in P^{(p,n)}$.

The identity (6.11) may be viewed as an A_{n-1} version of the second Rogers–Ramanujan identity and its higher moduli generalizations [1, 9]. By standard Bailey chain type arguments [3, 4, 53] it follows that for $k \geq 1$ and $a \in \{0, 1\}$,

$$X_{(12L)}^{2k+a,2}(q) = \sum_{L \geq n_1 \geq \dots \geq n_{k-1} \geq 0} \frac{q^{L+n_1(n_1+1)+\dots+n_{k-1}(n_{k-1}+1)} (q)_{2L}}{(q)_{L-n_1} \cdots (q)_{n_{k-2}-n_{k-1}} (q^{2-a}; q^{2-a})_{n_{k-1}}}$$

which substituted into (6.11) gives

$$(6.12) \quad \sum_{n_1 \geq \dots \geq n_k \geq 0} \frac{q^{n_1(n_1+1)+\dots+n_k(n_k+1)}}{(q)_{n_1-n_2} \cdots (q)_{n_{k-1}-n_k} (q^{2-a}; q^{2-a})_{n_k}} = \frac{(q, q^{2k+a+1}, q^{2k+a+2}; q^{2k+a+2})_\infty}{(q)_\infty}.$$

There is in fact an alternative route from (6.11) to this result, based on level-rank duality. First observe that if we write the right-hand side of (6.11) as $f_{n,p}(q)$, then $f_{n,p} = f_{p,n}$. The identities of (6.12) should therefore also result from the $p = 2$ case of (6.11). Indeed, experiments for $n = 3$ and 4 confirm the following conjecture:

$$X_\eta^{2,2k+a}(q) = \frac{q^{n_1+\dots+n_k} (q)_{m(\eta)}}{(q)_{n_1-n_2} \cdots (q)_{n_{k-1}-n_k} (q^{2-a}; q^{2-a})_{n_k}}$$

if $m(\eta) = (n_1 - n_2, \dots, n_{k-1} - n_k, n_k, n_k, n_{k-1} - n_k, \dots, n_1 - n_2)$ and zero otherwise. Here n_k, n_k should read $2n_k$ when $a = 0$. Substituted into (6.11) this once again yields (6.12).

As a final application we employ the following conjecture for $n = 3$ and $\Lambda \in P_+^{(2,4)}$:

$$X_{\eta, \Lambda, -\Lambda_0}(q) = q^{\frac{1}{6}(\zeta_1 - \zeta_2)^2 + \frac{1}{2}(\zeta_1 + \zeta_2) - \|\bar{\Lambda}\|^2} = q^{(\zeta_1^2 + 1) + (\zeta_2^2 + 1) - \frac{1}{2}\zeta C^{-1}\zeta - \|\bar{\Lambda}\|^2},$$

where $\zeta = m(\eta) \in \mathbb{Z}_+^2$. Since the elements of $P_+^{(2,4)}$ are given by $\Lambda_l - 2\Lambda_0$ for $l \in \{0, 1, 2\}$, the condition $|\eta| \bar{\Lambda}_1 - \bar{\Lambda} \in Q$ is equivalent to $\zeta_1 + 2\zeta_2 + l \equiv 0 \pmod{3}$. Substituting this last conjecture into (6.10) gives

$$\begin{aligned} & \frac{1}{(q)_\infty^2} \sum_{k \in Q} q^{12(k|k) + 6(k|\bar{\Lambda}_1) + 2(k|\rho)} \prod_{1 \leq i < j \leq 3} (1 - q^{4(k_i - k_j) + j - i + (\bar{\Lambda}_1|\varepsilon_i - \varepsilon_j)}) \\ & = \sum_{\substack{\zeta_1, \zeta_2 \geq 0 \\ \zeta_1 + 2\zeta_2 \equiv -l \pmod{3}}} \frac{q^{(\zeta_1^2 + 1) + (\zeta_2^2 + 1) - \frac{3}{2}\|\bar{\Lambda}_l\|^2}}{(q)_{\zeta_1} (q)_{\zeta_2}} \end{aligned}$$

whereas substitution into (6.3) leads to

$$b_{\Lambda_l - 2\Lambda_0, \Lambda_l - 2\Lambda_0, -2\Lambda_0}(q) = \sum_{\substack{\zeta_1, \zeta_2 \geq 0 \\ \zeta_1 + 2\zeta_2 \equiv -l \pmod{3}}} \frac{q^{\frac{1}{3}(\zeta_1 - \zeta_2)^2 + \zeta_1 + \zeta_2 - 2\|\bar{\Lambda}_l\|^2}}{(q)_{\zeta_1} (q)_{\zeta_2}}.$$

Acknowledgement. I thank Will Orrick, Anne Schilling and Trevor Welsh for many helpful discussions.

REFERENCES

- [1] G. E. Andrews, *An analytic generalization of the Rogers–Ramanujan identities for odd moduli*, *Prod. Nat. Acad. Sci. USA* **71** (1974), 4082–4085. 6
- [2] G. E. Andrews, *The Theory of Partitions*, *Encyclopedia of Mathematics and its Applications*, Vol. 2, (Addison-Wesley, Reading, Massachusetts, 1976). 2.3
- [3] G. E. Andrews, *Multiple series Rogers–Ramanujan type identities*, *Pacific J. Math.* **114** (1984), 267–283. 6
- [4] G. E. Andrews, *Bailey’s transform, lemma, chains and tree*, in *Special Functions 2000: Current Perspective and Future Directions*, pp. 1–22, J. Bustoz *et al.* eds., (Kluwer Academic Publishers, Dordrecht, 2001). 6
- [5] G. E. Andrews, A. Schilling and S. O. Warnaar, *An A_2 Bailey lemma and Rogers–Ramanujan-type identities*, *J. Amer. Math. Soc.* **12** (1999), 677–702. 1
- [6] W. N. Bailey, *Some identities in combinatorial analysis*, *Proc. London Math. Soc. (2)* **49** (1947), 421–435. 1
- [7] W. N. Bailey, *Identities of the Rogers–Ramanujan type*, *Proc. London Math. Soc. (2)* **50** (1949), 1–10. 1
- [8] A. Berkovich, *Fermionic counting of RSOS-states and Virasoro character formulas for the unitary minimal series $M(\nu, \nu + 1)$. Exact results*, *Nucl. Phys. B* **431** (1994), 315–348. 6
- [9] D. M. Bressoud, *An analytic generalization of the Rogers–Ramanujan identities with interpretation*, *Quart. J. Maths. Oxford (2)* **31** (1980), 385–399. 6
- [10] L. M. Butler, *Subgroup lattices and symmetric functions*, *Memoirs of the Amer. Math. Soc.*, no. 539, Vol. 112 (1994). 2.1
- [11] B. Feigin and A. V. Stoyanovsky, *Quasi-particle models for the representations of Lie algebras and geometry of flag manifold*, *Funct. Anal. Appl.* **28** (1994), 68–90. 4
- [12] W. Fulton, *Young tableaux: with applications to representation theory and geometry*, *London Math. Soc. student texts* **35**, (Cambridge University Press, Cambridge, 1997). 2.1
- [13] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, *Encyclopedia of Mathematics and its Applications*, Vol. 35, (Cambridge University Press, Cambridge, 1990). 1, 1, 2.3
- [14] G. Georgiev, *Combinatorial construction of modules for infinite-dimensional Lie algebras. II. Parafermionic space*, arXiv:q-alg/9504024. 4
- [15] G. Hatayama, A. N. Kirillov, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, *Character formulae of \widehat{sl}_n -modules and inhomogeneous paths*, *Nucl. Phys. B* **536** [PM] (1999), 575–616. 2.2, 3.1, 3.2, 3.3, 3.3, 3.4, 4
- [16] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, *Remarks on fermionic formula*, in *Recent Developments in Quantum Affine Algebras and Related Topics*, pp. 243–291, N. Jing and K. C. Misra eds., *Contemp. Math.* Vol. 248 (AMS, Providence, 1999). 3.1
- [17] V. G. Kac, *Infinite-dimensional Lie algebras, and the Dedekind η -function*, *Functional Anal. Appl.* **8** (1974), 68–70. 4
- [18] V. G. Kac, *Infinite-dimensional algebras, Dedekind’s η -function, classical Möbius function and the very strange formula*, *Adv. in Math.* **30** (1978), 85–136. 4
- [19] V. G. Kac, *An elucidation of: “Infinite-dimensional algebras, Dedekind’s η -function, classical Möbius function and the very strange formula”*. $E_8^{(1)}$ and the cube root of the modular invariant j , *Adv. in Math.* **35** (1980), 264–273. 4
- [20] V. G. Kac, *Infinite-dimensional Lie algebras*, third edition, (Cambridge University Press, Cambridge, 1990). 4, 4
- [21] V. G. Kac and D. H. Peterson, *Infinite-dimensional Lie algebras, theta functions and modular forms*, *Adv. in Math.* **53** (1984), 125–264. 4
- [22] V. G. Kac and M. Wakimoto, *Modular invariant representations of infinite-dimensional Lie algebras and superalgebras*, *Proc. Nat. Acad. Sci. USA* **85** (1988), 4956–4960. 5, 5
- [23] V. G. Kac and M. Wakimoto, *Branching functions for winding subalgebras and tensor products*, *Acta Appl. Math.* **21** (1990), 3–39. 5, 6, 6
- [24] A. N. Kirillov and N. Yu. Reshetikhin, *The Bethe Ansatz and the combinatorics of Young tableaux*, *J. Soviet Math.* **41** (1988), 925–955. 3.3

- [25] A. N. Kirillov, A. Schilling and M. Shimozono, *A bijection between Littlewood–Richardson tableaux and rigged configurations*, *Selecta Math. (N.S.)* **8** (2002), 67–135. 3.3
- [26] A. N. Kirillov and M. Shimozono, *A generalization of the Kostka–Foulkes polynomials*, *J. Algebraic Combin.* **15** (2002), 27–69. 2.2, 2.2, 3.3
- [27] A. Kuniba, T. Nakanishi and J. Suzuki, *Characters of conformal field theories from thermodynamic Bethe ansatz*, *Mod. Phys. Lett. A* **8** (1993) 1649–1660. 4
- [28] A. Lascoux and M.-P. Schützenberger, *Le monoïde plaxique*, in *Noncommutative Structures in Algebra and Geometric Combinatorics*, pp. 129–156, A. de Luca ed., *Quad. Ricerca Sci.* **109**, (CNR, Rome, 1981). 2.1
- [29] J. Lepowsky and M. Primc, *Structure of the standard modules for the affine Lie algebra $A_1^{(1)}$* , *Contemp. Math.* Vol. 46 (AMS, Providence, 1985). 4
- [30] D. E. Littlewood and A. R. Richardson, *Group characters and algebra*, *Phil. Trans. Royal. Soc. London. Ser. A* **233** (1934), 99–141. 2.2
- [31] I. G. Macdonald, *Affine root systems and Dedekind’s η -function*, *Invent. Math.* **15** (1972), 91–143. 6
- [32] I. G. Macdonald, *Symmetric functions and Hall polynomials*, second edition, (Oxford University Press, Oxford, 1995). 1, 1, 2.1, 2.2, 2.3
- [33] L. Manivel, *Symmetric functions, Schubert polynomials and degeneracy loci*, *SMF/AMS Texts and Monographs*, Vol. 6 (2001). 2.1
- [34] S. C. Milne, *Balanced ${}_3\phi_2$ summation theorems for $U(n)$ basic hypergeometric series*, *Adv. Math.* **131** (1997), 93–187. 1
- [35] S. C. Milne, *Transformations of $U(n+1)$ multiple basic hypergeometric series*, in *Physics and Combinatorics*, pp. 201–243, A. N. Kirillov et al. eds., (World Scientific, Singapore, 2001). 1
- [36] S. C. Milne and G. M. Lilly, *The A_ℓ and C_ℓ Bailey transform and lemma*, *Bull. Amer. Math. Soc. (N.S.)* **26** (1992), 258–263. 1
- [37] S. C. Milne and G. M. Lilly, *Consequences of the A_ℓ and C_ℓ Bailey transform and Bailey lemma*, *Discrete Math.* **139** (1995), 319–346. 1
- [38] M. Okado, A. Schilling and M. Shimozono, *Crystal bases and q -identities*, in *q -Series with Applications to Combinatorics, Number Theory, and Physics*, pp. 29–53, B. C. Berndt and K. Ono eds., *Contemp. Math.* Vol. 291 (AMS, Providence, 2001). 4, 6
- [39] L. J. Rogers, *Second memoir on the expansion of certain infinite products*, *Proc. London Math. Soc.* **25** (1894), 318–343. 1
- [40] A. Schilling and M. Shimozono, *Bosonic formula for level-restricted paths*, in *Combinatorial Methods in Representation Theory*, pp. 305–325, K. Koike et al. eds., *Adv. Studies in Pure Math.* **28**, (Kinokuniya, Tokyo, 2000). 4, 4, 4, 6
- [41] A. Schilling and M. Shimozono, *Fermionic formulas for level-restricted generalized Kostka polynomials and coset branching functions*, *Commun. Math. Phys.* **220** (2001), 105–164. 3.1, 4, 6
- [42] A. Schilling and S. O. Warnaar, *A higher level Bailey lemma: proof and application*, *The Ramanujan Journal* **2** (1998), 327–349. 1, 3.1, 4
- [43] A. Schilling and S. O. Warnaar, *Supernomial coefficients, polynomial identities and q -series*, *The Ramanujan Journal* **2** (1998), 459–494. 2.2, 3.4
- [44] A. Schilling and S. O. Warnaar, *Inhomogeneous lattice paths, generalized Kostka polynomials and A_{n-1} supernomials*, *Commun. Math. Phys.* **202** (1999), 359–401. 1, 2.2, 2.2, 3.1, 3.3
- [45] A. Schilling and S. O. Warnaar, *A generalization of the q -Saalschütz sum and the Burge transform*, in *Physical Combinatorics*, pp. 163–183, M. Kashiwara and T. Miwa eds., *Progress in Mathematics* **191**, (Birkhäuser, Boston, 2000). 1
- [46] A. Schilling and S. O. Warnaar, *Conjugate Bailey pairs*, in *Recent Developments in Infinite-Dimensional Lie Algebras and Conformal Field Theory*, S. Berman et al. eds., *Contemp. Math.* Vol. 297 (AMS, Providence, 2002). 1, 5, 6
- [47] M. Shimozono, *A cyclage poset structure for Littlewood–Richardson tableaux*, *European J. Combin.* **22** (2001), 365–393. 2.2
- [48] M. Shimozono, *Multi-atoms and monotonicity of generalized Kostka polynomials*, *European J. Combin.* **22** (2001), 395–414. 2.2
- [49] M. Shimozono, *Affine type A crystal structure on tensor products of rectangles, Demazure characters, and nilpotent varieties*, *J. Algebraic Combin.* **15** (2002), 151–187. 2.2
- [50] M. Shimozono and J. Weyman, *Graded characters of modules supported in the closure of a nilpotent conjugacy class*, *European J. Combin.* **21** (2000), 257–288. 2.2
- [51] S. O. Warnaar, *Fermionic solution of the Andrews–Baxter–Forrester model. II. Proof of Melzer’s polynomial identities*, *J. Stat. Phys.* **84** (1996), 49–83. 6
- [52] S. O. Warnaar, *Supernomial coefficients, Bailey’s lemma and Rogers–Ramanujan-type identities. A survey of results and open problems*, *Sém. Lothar. Combin.* **42** (1999), Art. B42n, 22 pp. 1, 3.1

- [53] S. O. Warnaar, *50 Years of Bailey's lemma*, in *Algebraic Combinatorics and Applications*, pp. 333–347, A. Betten *et al.* eds., (Springer, Berlin, 2001). 6

DEPARTMENT OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF MELBOURNE, VIC 3010, AUSTRALIA